

WORLD  
INTELLECTUAL  
PROPERTY  
ORGANIZATION



IP SERVICES

Home IP Services PatentScope Patent Search



Search result: 1 of 1

## (WO/2002/020546) ANTISENSE MODULATION OF PHOSPHORYLASE KINASE ALPHA EXPRESSION

Biblio. Data Description Claims National Phase Notices Documents

Latest bibliographic data on file with the International Bureau

Publication Number: WO/2002/020546 International Application No.: PCT/US2001/026608  
Publication Date: 14.03.2002 International Filing Date: 24.08.2001  
Chapter 2 Demand Filed: 27.03.2002

Int. Class.: A61K 38/00 (2006.01), A61K 48/00 (2006.01), C12N 15/11 (2006.01)

Applicants: ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue Carlsbad, CA 92008 (US).  
MONIA, Brett, P. [US/US]; 7605 Neuva Castilla Way La Costa, CA 92009 (US) (US Only).  
WYATT, Jacqueline, R. [US/US]; 1065 Hymettus Avenue Encinitas, CA 92024 (US) (US Only).

Inventors: MONIA, Brett, P. [US/US]; 7605 Neuva Castilla Way La Costa, CA 92009 (US).  
WYATT, Jacqueline, R. [US/US]; 1065 Hymettus Avenue Encinitas, CA 92024 (US).

Agent: LICATA, Jane, Massey; Licata & Tyrrell P.C. 66 E. Main Street Marlton, NJ 08053 (US).

Priority Data: 09/657,452 07.09.2000 US

Title: ANTISENSE MODULATION OF PHOSPHORYLASE KINASE ALPHA 1 EXPRESSION

Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of Phosphorylase kinase alpha 1. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding Phosphorylase kinase alpha 1. Methods of using these compounds for modulation of Phosphorylase kinase alpha 1 expression and for treatment of diseases associated with expression of Phosphorylase kinase alpha 1 are provided.

Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, D, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, T, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.  
African Regional Intellectual Property Org. (ARIPO) (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW)  
Eurasian Patent Organization (EAPO) (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM)  
European Patent Office (EPO) (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR)  
African Intellectual Property Organization (OAPI) (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Publication Language:

English (EN)

Filing Language:

English (EN)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number  
**WO 02/20546 A1**

- (51) International Patent Classification<sup>7</sup>: **C07H 21/04**, (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, ZA, ZW.
- A61K 48/00, C12Q 1/68, C12N 15/63
- (21) International Application Number: PCT/US01/26608.
- (22) International Filing Date: 24 August 2001 (24.08.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/657,452 7 September 2000 (07.09.2000) US
- (71) Applicant (*for all designated States except US*): **ISIS PHARMACEUTICALS, INC.** [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **MONIA, Brett, P.** [US/US]; 7605 Neuva Castilla Way, La Costa, CA 92009 (US). **WYATT, Jacqueline, R.** [US/US]; 1065 Hymettus Avenue, Encinitas, CA 92024 (US).
- (74) Agent: **LICATA, Jane, Massey**; Licata & Tyrrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**WO 02/20546 A1**

(54) Title: ANTISENSE MODULATION OF PHOSPHORYLASE KINASE ALPHA 1 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of Phosphorylase kinase alpha 1. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding Phosphorylase kinase alpha 1. Methods of using these compounds for modulation of Phosphorylase kinase alpha 1 expression and for treatment of diseases associated with expression of Phosphorylase kinase alpha 1 are provided.

**ANTISENSE MODULATION OF  
PHOSPHORYLASE KINASE ALPHA 1 EXPRESSION**

**5 FIELD OF THE INVENTION**

The present invention provides compositions and methods for modulating the expression of Phosphorylase kinase alpha 1. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with  
10 nucleic acids encoding Phosphorylase kinase alpha 1. Such compounds have been shown to modulate the expression of Phosphorylase kinase alpha 1.

**BACKGROUND OF THE INVENTION**

Balanced energy metabolism is critical to the regulation  
15 of all biological processes. In higher organisms, energy stores are in the form of glycogen. Upon energy deficit, these stores are mobilized through enzymatic digestion to glucose-1-phosphate by a diverse set of signals and are used to maintain blood-glucose levels, as a source of energy  
20 during muscle contraction and as source of fuel for a broad range of cellular activities.

The protein kinase, phosphorylase kinase (PHK) plays a central role in the regulation of glycogen degradation or glycogenolysis by phosphorylating glycogen phosphorylase b, a  
25 unique reaction catalyzed only by phosphorylase kinase. It also lies at the interface between signaling and metabolic pathways and translates the pleiotropic actions of extracellular signals, including hormonal and neuronal, into specific and directional intracellular responses. In  
30 addition, phosphorylase kinase can express varying degrees of activity depending on pH, metal ion concentration, allosteric effectors and covalent modifications (Brushia and Walsh, *Front. Biosci.*, 1999, 4, D618-641).

Structurally, phosphorylase kinase is one of the most

complex enzymes isolated to date, a hexadecamer, having three distinct regulatory subunits, alpha, beta and delta (also known as calmodulin), and one catalytic subunit, gamma. Each holoenzyme is composed of four heterotetramers of the  
5 component subunits and the subunit stoichiometry has been shown to vary depending on the tissue source. The phosphorylase kinase subunits also exist as multiple isoforms adding an additional layer of complexity. The alpha, beta, and gamma isoforms are found expressed in the liver and  
10 muscle with minor amounts in the gut, while the delta (calmodulin) isoforms are expressed in all tissues examined (Brushia and Walsh, *Front. Biosci.*, 1999, 4, D618-641).

Due to the direct relationship between phosphorylase kinase enzyme activity and maintenance of blood-glucose  
15 homeostasis, modifications to the regulatory properties of this enzyme could provide great therapeutic benefit in the arena of metabolic disorders, especially diabetes.

Phosphorylase kinase alpha 1 (also known as PHKA, PHKA1 and  $\alpha M$ ) is one of the three regulatory alpha subunit isoforms  
20 identified to date and is localized solely in muscle tissue (Wullrich et al., *J. Biol. Chem.*, 1993, 268, 23208-23214).

The gene is located on chromosome Xq13 and several types of mutations in this gene have been reported which result in differential mRNA processing and certain forms of glycogen  
25 storage diseases. These mutations include a splice junction mutation in a patient with myopathy (Bruno et al., *Biochem. Biophys. Res. Commun.*, 1998, 249, 648-651) and nonsense mutations that resemble the X-linked phosphorylase kinase deficiency seen in I-strain mice (Wehner et al., *Hum. Mol.*  
30 *Genet.*, 1994, 3, 1983-1987). In mice this nonsense mutation results in a frameshift of the coding region and therefore disrupts the expression of both the liver and muscle isoforms of the alpha subunit (Bender, *Biochem. Biophys. Res. Commun.*, 1991, 179, 707-712; Bender and Lalley, *Proc. Natl. Acad. Sci.*

U. S. A., 1989, 86, 9996-10000; Schneider et al., Nat. Genet., 1993, 5, 381-385).

Currently however, there are no known therapeutic agents which effectively inhibit the synthesis of phosphorylase kinase alpha 1 and to date, investigative strategies aimed at studying phosphorylase kinase alpha 1 function have involved the use of antibodies and crosslinking agents.

Consequently, there remains a long felt need for agents capable of effectively modulating phosphorylase kinase alpha 1 function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of phosphorylase kinase alpha 1 expression.

The present invention provides compositions and methods for modulating phosphorylase kinase alpha 1 expression.

#### SUMMARY OF THE INVENTION

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding Phosphorylase kinase alpha 1, and which modulate the expression of Phosphorylase kinase alpha 1. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of Phosphorylase kinase alpha 1 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of Phosphorylase kinase alpha 1 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the

invention.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric compounds, particularly antisense oligonucleotides, for use in  
5 modulating the function of nucleic acid molecules encoding Phosphorylase kinase alpha 1, ultimately modulating the amount of Phosphorylase kinase alpha 1 produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids  
10 encoding Phosphorylase kinase alpha 1. As used herein, the terms "target nucleic acid" and "nucleic acid encoding Phosphorylase kinase alpha 1" encompass DNA encoding Phosphorylase kinase alpha 1, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from  
15 such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as  
20 "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of  
25 the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of Phosphorylase kinase alpha 1. In the context of the present  
30 invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

-5-

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the  
5 identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the  
10 present invention, the target is a nucleic acid molecule encoding Phosphorylase kinase alpha 1. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression  
15 of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is  
20 typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG,  
25 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or  
30 formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of

-6-

conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding Phosphorylase kinase alpha 1, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an



N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain

position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that  
5 position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms  
10 which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100%  
15 complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility,  
20 and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of  
25 in vitro assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of  
30 these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as "active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the

invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense  
5 oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a  
10 biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic  
15 moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic  
20 modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of  
25 ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions  
30 which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The  
5 antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from  
10 about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

15 As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate  
20 group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently  
25 link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure,  
30 the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds

-11-

useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus  
5 atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to  
10 be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-  
15 alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and borano-  
20 phosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most  
25 internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the  
30 preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925;

5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253;  
5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899;  
5,721,218; 5,672,697 and 5,625,050, certain of which are  
commonly owned with this application, and each of which is  
5 herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not  
include a phosphorus atom therein have backbones that are  
formed by short chain alkyl or cycloalkyl internucleoside  
linkages, mixed heteroatom and alkyl or cycloalkyl  
10 internucleoside linkages, or one or more short chain  
heteroatomic or heterocyclic internucleoside linkages. These  
include those having morpholino linkages (formed in part from  
the sugar portion of a nucleoside); siloxane backbones;  
sulfide, sulfoxide and sulfone backbones; formacetyl and  
15 thioformacetyl backbones; methylene formacetyl and  
thioformacetyl backbones; riboacetyl backbones; alkene  
containing backbones; sulfamate backbones; methyleneimino and  
methylenediazino backbones; sulfonate and sulfonamide  
backbones; amide backbones; and others having mixed N, O, S  
20 and CH<sub>2</sub> component parts.

Representative United States patents that teach the  
preparation of the above oligonucleosides include, but are  
not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444;  
5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564;  
25 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677;  
5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289;  
5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070;  
5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and  
5,677,439, certain of which are commonly owned with this  
30 application, and each of which is herein incorporated by  
reference.

In other preferred oligonucleotide mimetics, both the  
sugar and the internucleoside linkage, i.e., the backbone, of  
the nucleotide units are replaced with novel groups. The

-13-

base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylinino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are

-14-

$O[(CH_2)_nO]_mCH_3$ ,  $O(CH_2)_nOCH_3$ ,  $O(CH_2)_nNH_2$ ,  $O(CH_2)_nCH_3$ ,  $O(CH_2)_nONH_2$ ,  
and  $O(CH_2)_nON[(CH_2)_nCH_3]_2$ , where n and m are from 1 to about  
10. Other preferred oligonucleotides comprise one of the  
following at the 2' position:  $C_1$  to  $C_{10}$  lower alkyl,  
5 substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl,  
O-alkaryl or O-aralkyl, SH,  $SCH_3$ , OCN, Cl, Br, CN,  $CF_3$ ,  $OCF_3$ ,  
 $SOCH_3$ ,  $SO_2CH_3$ ,  $ONO_2$ ,  $NO_2$ ,  $N_3$ ,  $NH_2$ , heterocycloalkyl,  
heterocycloalkaryl, aminoalkylamino, polyalkylamino,  
substituted silyl, an RNA cleaving group, a reporter group,  
10 an intercalator, a group for improving the pharmacokinetic  
properties of an oligonucleotide, or a group for improving  
the pharmacodynamic properties of an oligonucleotide, and  
other substituents having similar properties. A preferred  
modification includes 2'-methoxyethoxy (2'-O- $CH_2CH_2OCH_3$ , also  
15 known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al.,  
*Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy  
group. A further preferred modification includes 2'-  
dimethylaminoethoxy, i.e., a  $O(CH_2)_2ON(CH_3)_2$  group, also  
known as 2'-DMAOE, as described in examples hereinbelow, and  
20 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-  
dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O- $CH_2-O-CH_2-$   
 $N(CH_3)_2$ , also described in examples hereinbelow.-

A further preferred modification includes Locked Nucleic  
Acids (LNAs) in which the 2'-hydroxyl group is linked to the  
25 3' or 4' carbon atom of the sugar ring thereby forming a  
bicyclic sugar moiety. The linkage is preferably a methylene  
( $-CH_2-$ )<sub>n</sub> group bridging the 2' oxygen atom and the 3' or 4'  
carbon atom wherein n is 1 or 2. LNAs and preparation thereof  
are described in WO 98/39352 and WO 99/14226.

30 Other preferred modifications include 2'-methoxy (2'-O-  
 $CH_3$ ), 2'-aminopropoxy (2'-O- $CH_2CH_2CH_2NH_2$ ), 2'-allyl (2'- $CH_2-$   
 $CH=CH_2$ ), 2'-O-alkyl (2'-O- $CH_2-CH=CH_2$ ) and 2'-fluoro (2'-F).  
The 2'-modification may be in the arabino (up) position or  
ribo (down) position. A preferred 2'-arabino modification is



2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 10 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of 15 which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine 20 (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine 25 and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ( $-C\equiv C-CH_3$ ) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 30 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and

-16-

8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one),  
5 phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified  
10 nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed  
15 in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by  
20 Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-  
25 substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and  
30 Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the

-17-

preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein  
10 incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide  
15 one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of  
20 the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugates groups include  
25 cholesterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake,  
30 enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative

-18-

conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid

5 moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et

10 al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330;

15 Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a

20 polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or

25 hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+) -

30 pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an

antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

5 Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802;  
10 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506;  
15 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the  
20 instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a  
25 single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense compounds, particularly  
30 oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to

-20-

confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a  
5 substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the  
10 efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA  
15 target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more  
20 oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not  
25 limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

30 The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other

means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

5       The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules. The compounds of the invention may also be admixed,  
10 encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative  
15 United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556;  
20 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

25       The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite  
30 or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is

-22-

prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the

5 oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphatel] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach et al.

10 The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

15 Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are

20 N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds are

25 prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms

30 differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically



acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and  
5 phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or  
10 phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid,  
15 gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20  
20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfoic acid,  
25 naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be  
30 prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of Phosphorylase kinase alpha 1 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding Phosphorylase kinase alpha 1, enabling sandwich and other assays to easily be constructed

to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding Phosphorylase kinase alpha 1 can be detected by means known in the art. Such means may include conjugation  
5 of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of Phosphorylase kinase alpha 1 in a sample may also be prepared.

10 The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is  
15 desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and  
20 transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-  
25 methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids  
30 and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in

-26-

admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE  
5 ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may  
10 be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic  
15 acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g.  
20 isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

25 Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents,  
30 emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include

fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate, . Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate),

poly(butylcyanoacrylate), poly(isobutylcyanoacrylate),  
poly(isohexylcyanoacrylate), DEAE-methacrylate, DEAE-  
hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-  
dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-  
5 lactic acid), poly(DL-lactic-co-glycolic acid (PLGA),  
alginate, and polyethyleneglycol (PEG). Oral formulations  
for oligonucleotides and their preparation are described in  
detail in United States applications 08/886,829 (filed July  
1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed  
10 February 23, 1999), 09/082,624 (filed May 21, 1998) and  
09/315,298 (filed May 20, 1999) each of which is incorporated  
herein by reference in their entirety.

Compositions and formulations for parenteral,  
intrathecal or intraventricular administration may include  
15 sterile aqueous solutions which may also contain buffers,  
diluent and other suitable additives such as, but not  
limited to, penetration enhancers, carrier compounds and  
other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention  
20 include, but are not limited to, solutions, emulsions, and  
liposome-containing formulations. These compositions may be  
generated from a variety of components that include, but are  
not limited to, preformed liquids, self-emulsifying solids  
and self-emulsifying semisolids.

25 The pharmaceutical formulations of the present  
invention, which may conveniently be presented in unit dosage  
form, may be prepared according to conventional techniques  
well known in the pharmaceutical industry. Such techniques  
include the step of bringing into association the active  
30 ingredients with the pharmaceutical carrier(s) or  
excipient(s). In general the formulations are prepared by  
uniformly and intimately bringing into association the active  
ingredients with liquid carriers or finely divided solid  
carriers or both, and then, if necessary, shaping the

product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid  
5 syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for  
10 example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as,  
15 but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the  
20 pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are  
25 typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical*  
30 *Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical*

Sciences, Mack Publishing Co., Easton, PA, 1985, p. 301).  
Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil  
5 (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily  
10 phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase.  
15 Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and  
20 water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a  
25 system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into  
30 the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing



emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption  
5 bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of  
10 emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York,  
15 N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in  
20 categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker  
25 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties  
30 such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations.

These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments  
5 and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants,  
10 hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New  
15 York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for  
20 example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-  
25 phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these  
30 formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion

formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic  
5 acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature  
10 (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability  
15 standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1,  
20 p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are  
25 formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New  
30 York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system.

Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in:  
5 *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the  
10 microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*,  
15 Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*,  
20 Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions  
25 offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic  
30 surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate

(MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to  
5 increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying  
10 microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to,  
15 materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone  
20 oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs,  
25 including peptides (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug  
30 absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm.*

*Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or  
5 oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic  
10 absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

15 Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of  
20 the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic*  
25 *Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the  
30 formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term

-37-

"liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

10 In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through  
15 such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect  
20 encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations  
25 are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological  
30 membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).



One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A

into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside  $G_{M1}$ , or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al., *Cancer Research*, 1993, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside  $G_{M1}$ , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside  $G_{M1}$  or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum et al. (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klivanov et al. (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to  
5 Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes  
10 comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be  
15 described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin),  
20 self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the  
25 skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The  
30 most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing

the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is  
5 classified as a nonionic surfactant. Nonionic surfactants  
find wide application in pharmaceutical and cosmetic products  
and are usable over a wide range of pH values. In general  
their HLB values range from 2 to about 18 depending on their  
structure. Nonionic surfactants include nonionic esters such  
10 as ethylene glycol esters, propylene glycol esters, glyceryl  
esters, polyglyceryl esters, sorbitan esters, sucrose esters,  
and ethoxylated esters. Nonionic alkanolamides and ethers  
such as fatty alcohol ethoxylates, propoxylated alcohols, and  
ethoxylated/propoxylated block polymers are also included in  
15 this class. The polyoxyethylene surfactants are the most  
popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge  
when it is dissolved or dispersed in water, the surfactant is  
classified as anionic. Anionic surfactants include  
20 carboxylates such as soaps, acyl lactylates, acyl amides of  
amino acids, esters of sulfuric acid such as alkyl sulfates  
and ethoxylated alkyl sulfates, sulfonates such as alkyl  
benzene sulfonates, acyl isethionates, acyl taurates and  
sulfosuccinates, and phosphates. The most important members  
25 of the anionic surfactant class are the alkyl sulfates and  
the soaps.

If the surfactant molecule carries a positive charge  
when it is dissolved or dispersed in water, the surfactant is  
classified as cationic. Cationic surfactants include  
30 quaternary ammonium salts and ethoxylated amines. The  
quaternary ammonium salts are the most used members of this  
class.

If the surfactant molecule has the ability to carry  
either a positive or negative charge, the surfactant is

-44-

classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example,

sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al.,  
5 *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid,  
10 linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl),  
15 and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et  
20 al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman  
25 et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935).

Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts  
30 of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid

(sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).



Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity

by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate,

sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic  
5 excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols,  
10 polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-  
15 aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral  
20 administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium  
25 stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally  
30 found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or

anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-

hydroxyperoxycyclophosphoramide, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, 5 cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), 10 sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not 15 limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th 20 Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

25 In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of 30 antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and

responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be  
5 calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can  
10 generally be estimated based on  $EC_{50}$ s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art  
15 can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state,  
20 wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred  
25 embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

#### EXAMPLES

##### Example 1

##### Nucleoside Phosphoramidites for Oligonucleotide Synthesis 30 Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are

-53-

prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

#### **2'-Fluoro amidites**

##### **2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a  $S_N2$ -displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was, selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

##### **2'-Fluorodeoxyguanosine**

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group

was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then  
5 deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

#### **2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-  
10 anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination  
15 of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-O-(2-Methoxyethyl) modified amidites**

20 2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

#### **2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]**

25 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide  
30 gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum



-55-

amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

**2'-O-Methoxyethyl-5-methyluridine**

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH<sub>3</sub>CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH<sub>2</sub>Cl<sub>2</sub>/acetone/MeOH (20:5:3) containing 0.5% Et<sub>3</sub>NH. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was

-56-

added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH<sub>3</sub>CN (200 mL).

5 The residue was dissolved in CHCl<sub>3</sub> (1.5 L) and extracted with 2x500 mL of saturated NaHCO<sub>3</sub> and 2x500 mL of saturated NaCl. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted  
10 with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et<sub>3</sub>NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

15       3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and  
20 stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl<sub>3</sub> (800 mL) and  
25 extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl<sub>3</sub>. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a  
30 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

**3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine**

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH<sub>3</sub>CN (700 mL) and set aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH<sub>3</sub>CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl<sub>3</sub> was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO<sub>3</sub> and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

**2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH<sub>4</sub>OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH<sub>3</sub> gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl<sub>3</sub> (700 mL) and extracted with saturated NaHCO<sub>3</sub> (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO<sub>4</sub> and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et<sub>3</sub>NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

**N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite**

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra-(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated NaHCO<sub>3</sub> (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (300 mL), and the extracts were combined, dried over MgSO<sub>4</sub> and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also  
5 known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside  
amidites] are prepared as described in the following  
paragraphs. Adenosine, cytidine and guanosine nucleoside  
amidites are prepared similarly to the thymidine (5-  
methyluridine) except the exocyclic amines are protected with  
10 a benzoyl moiety in the case of adenosine and cytidine and  
with isobutyl in the case of guanosine.

5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-  
methyluridine

O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese,  
15 Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g,  
0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml)  
at ambient temperature under an argon atmosphere and with  
mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g,  
119.0mL, 1.1eq, 0.458mmol) was added in one portion. The  
20 reaction was stirred for 16 h at ambient temperature. TLC  
(R<sub>f</sub> 0.22, ethyl acetate) indicated a complete reaction. The  
solution was concentrated under reduced pressure to a thick  
oil. This was partitioned between dichloromethane (1 L) and  
saturated sodium bicarbonate (2x1 L) and brine (1 L). The  
25 organic layer was dried over sodium sulfate and concentrated  
under reduced pressure to a thick oil. The oil was dissolved  
in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and  
the solution was cooled to  
-10°C. The resulting crystalline product was collected by  
30 filtration, washed with ethyl ether (3x200 mL) and dried  
(40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and  
NMR were consistent with pure product.

-60-

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-2'-O-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (R<sub>f</sub> 0.67 for desired product and R<sub>f</sub> 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with

-61-

triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over  $P_2O_5$  under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, 5 sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition 10 was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-15 phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

**5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine**

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-20 methyluridine (3.1g, 4.5mmol) was dissolved in dry  $CH_2Cl_2$  (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold  $CH_2Cl_2$ , and the combined organic phase was washed with water, brine and dried over 25 anhydrous  $Na_2SO_4$ . The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue 30 chromatographed to get 5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

-62-

**5'-O-tert-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine**

5'-O-tert-butyldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was  
5 dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the  
10 ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, evaporated to dryness. Residue was  
15 dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and  
20 reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO<sub>3</sub> (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over  
25 anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

30      **2'-O-(dimethylaminoxyethyl)-5-methyluridine**

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-



-63-

dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Solvent was removed under vacuum and the residue placed on a flash column and  
5 eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine 2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at  
10 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture  
15 was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

20 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To  
25 the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added.  
30 The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). Ethyl

-64-

acetate layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

**N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-

diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-  
5 O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50  
10 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O<sup>2</sup>-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed  
15 in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with  
20 ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are  
25 concentrated a colorless solid forms which is collected to give the title compound as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL),  
30 triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers are washed

-66-

with saturated  $\text{NaHCO}_3$  solution, followed by saturated  $\text{NaCl}$  solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using  $\text{MeOH}:\text{CH}_2\text{Cl}_2:\text{Et}_3\text{N}$  (20:1, v/v, with 1% triethylamine) gives the title compound.

5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in  $\text{CH}_2\text{Cl}_2$  (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

#### Example 2

##### Oligonucleotide synthesis

20 Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at  $55^\circ\text{C}$  (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M  $\text{NaCl}$  solution. Phosphinate oligonucleotides are prepared as described in

U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

5        3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878,  
10 herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

15        3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by  
20 reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### 25 Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked  
30 oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are

prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides  
5 are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

#### 10 Example 4

##### PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential  
15 Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

##### Example 5

#### 20 Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between  
25 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of  
30 the second type are also known in the art as "hemimers" or "wingmers".

[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric

Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl

-69-

phosphorothioate and 2'-deoxy phosphorothioate oligo-nucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated  
5 synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphor-amidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s  
10 repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room  
15 temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac  
20 before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-  
25 (Methoxyethyl)] Chimeric Phosphorothioate  
Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxy-ethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl  
30 chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

[2'-O-(2-Methoxyethyl)Phosphodiester] -- [2'-deoxy  
Phosphorothioate] -- [2'-O-(2-Methoxyethyl)  
Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester) -- [2'-deoxy phos-  
5 phorothioate] -- [2'-O-(methoxyethyl) phosphodiester] chimeric  
oligonucleotides are prepared as per the above procedure for  
the 2'-O-methyl chimeric oligonucleotide with the  
substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-  
methyl amidites, oxidization with iodine to generate the  
10 phosphodiester internucleotide linkages within the wing  
portions of the chimeric structures and sulfurization  
utilizing 3,4-dihydro-1,2-benzodithiole-3-one 1,1-dioxide (Beaucage  
Reagent) to generate the phosphorothioate internucleotide  
linkages for the center gap.

15 Other chimeric oligonucleotides, chimeric oligonucleo-  
sides and mixed chimeric oligonucleotides/oligonucleosides  
are synthesized according to United States patent 5,623,065,  
herein incorporated by reference.

#### Example 6

#### 20 Oligonucleotide Isolation

After cleavage from the controlled pore glass column  
(Applied Biosystems) and deblocking in concentrated ammonium  
hydroxide at 55°C for 18 hours, the oligonucleotides or  
oligonucleosides are purified by precipitation twice out of  
25 0.5 M NaCl with 2.5 volumes ethanol. Synthesized  
oligonucleotides were analyzed by polyacrylamide gel  
electrophoresis on denaturing gels and judged to be at least  
85% full length material. The relative amounts of  
phosphorothioate and phosphodiester linkages obtained in  
30 synthesis were periodically checked by <sup>31</sup>P nuclear magnetic  
resonance spectroscopy, and for some studies oligonucleotides  
were purified by HPLC, as described by Chiang et al., *J.*  
*Biol. Chem.* 1991, 266, 18162-18171. Results obtained with  
HPLC-purified material were similar to those obtained with



-71-

non-HPLC purified material.

**Example 7**

**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III)  
5 phosphoramidite chemistry on an automated synthesizer capable  
of assembling 96 sequences simultaneously in a standard 96  
well format. Phosphodiester internucleotide linkages were  
afforded by oxidation with aqueous iodine. Phosphorothioate  
internucleotide linkages were generated by sulfurization  
10 utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage  
Reagent) in anhydrous acetonitrile. Standard base-protected  
beta-cyanoethyl-diisopropyl phosphoramidites were purchased  
from commercial vendors (e.g. PE-Applied Biosystems, Foster  
City, CA, or Pharmacia, Piscataway, NJ). Non-standard  
15 nucleosides are synthesized as per known literature or  
patented methods. They are utilized as base protected beta-  
cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and  
deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature  
20 (55-60°C) for 12-16 hours and the released product then dried  
in vacuo. The dried product was then re-suspended in sterile  
water to afford a master plate from which all analytical and  
test plate samples are then diluted utilizing robotic  
pipettors.

**25 Example 8**

**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was  
assessed by dilution of samples and UV absorption  
spectroscopy. The full-length integrity of the individual  
30 products was evaluated by capillary electrophoresis (CE) in  
either the 96 well format (Beckman P/ACE™ MDQ) or, for  
individually prepared samples, on a commercial CE apparatus  
(e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone  
composition was confirmed by mass analysis of the compounds

utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were  
5 at least 85% full length.

#### Example 9

##### Cell culture and oligonucleotide treatment

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types  
10 provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 6 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is  
15 expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

##### T-24 cells:

20 The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum  
25 (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well  
30 plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium

-73-

and oligonucleotide.

A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

3T3-L1 cells:

The mouse embryonic adipocyte-like cell line 3T3-L1 was obtained from the American Type Culture Collection (Manassas, VA). 3T3-L1 cells were routinely cultured in DMEM, high glucose (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 80% confluence. Cells were seeded into 96-well plates (Falcon-

-74-

Primaria #3872) at a density of 4000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

A10 cells:

The rat aortic smooth muscle cell line A10 was obtained from the American Type Culture Collection (Manassas, VA). A10 cells were routinely cultured in DMEM, high glucose (American Type Culture Collection, Manassas, VA) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 80% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 2500 cells/well for use in RT-PCR analysis.

For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200  $\mu$ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130  $\mu$ L of OPTI-MEM™-1 containing 3.75  $\mu$ g/mL LIPOFECTIN™ (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive

control oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**,  
SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls  
shown in bold) with a phosphorothioate backbone which is  
targeted to human H-ras. For mouse or rat cells the positive  
5 control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCCAAGGA**,  
SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls  
shown in bold) with a phosphorothioate backbone which is  
targeted to both mouse and rat c-raf. The concentration of  
positive control oligonucleotide that results in 80%  
10 inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS  
15770) mRNA is then utilized as the screening concentration  
for new oligonucleotides in subsequent experiments for that  
cell line. If 80% inhibition is not achieved, the lowest  
concentration of positive control oligonucleotide that  
15 results in 60% inhibition of H-ras or c-raf mRNA is then  
utilized as the oligonucleotide screening concentration in  
subsequent experiments for that cell line. If 60% inhibition  
is not achieved, that particular cell line is deemed as  
unsuitable for oligonucleotide transfection experiments.

#### 20 Example 10

##### **Analysis of oligonucleotide inhibition of Phosphorylase kinase alpha 1 expression**

Antisense modulation of Phosphorylase kinase alpha 1  
expression can be assayed in a variety of ways known in the  
25 art. For example, Phosphorylase kinase alpha 1 mRNA levels  
can be quantitated by, e.g., Northern blot analysis,  
competitive polymerase chain reaction (PCR), or real-time PCR  
(RT-PCR). Real-time quantitative PCR is presently preferred.  
RNA analysis can be performed on total cellular RNA or  
30 poly(A)+ mRNA. Methods of RNA isolation are taught in, for  
example, Ausubel, F.M. et al., *Current Protocols in Molecular  
Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John  
Wiley & Sons, Inc., 1993. Northern blot analysis is routine  
in the art and is taught in, for example, Ausubel, F.M. et

al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

Protein levels of Phosphorylase kinase alpha 1 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to Phosphorylase kinase alpha 1 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991.

**Example 11****Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, 1996, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

**Example 12****Total RNA Isolation**

Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ L cold PBS. 100  $\mu$ L Buffer RLT was added to each well and the plate vigorously

-78-

agitated for 20 seconds. 100  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60  $\mu$ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60  $\mu$ L water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

### 25 Example 13

#### Real-time Quantitative PCR Analysis of Phosphorylase kinase alpha 1 mRNA Levels

Quantitation of Phosphorylase kinase alpha 1 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As



-79-

opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an

5 oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the

10 probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye.

15 During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the

20 remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics

25 built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test

30 samples.

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the

internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for  
5 GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of  
10 the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

15 PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25  $\mu$ L PCR cocktail (1x TAQMAN<sup>™</sup> buffer A, 5.5 mM MgCl<sub>2</sub>, 300  $\mu$ M each of dATP, dCTP and dGTP, 600  $\mu$ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase  
20 inhibitor, 1.25 Units AMPLITAQ GOLD<sup>™</sup>, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD<sup>™</sup>, 40 cycles  
25 of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene  
30 whose expression is constant, or by quantifying total RNA using RiboGreen<sup>™</sup> (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen<sup>™</sup> RNA quantification

-81-

reagent from Molecular Probes. Methods of RNA quantification by RiboGreen™ are taught in Jones, L.J., et al, *Analytical Biochemistry*, 1998, 265, 368-374.

In this assay, 175 µL of RiboGreen™ working reagent  
5 (RiboGreen™ reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25uL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

10 Probes and primers to human Phosphorylase kinase alpha 1 were designed to hybridize to a human Phosphorylase kinase alpha 1 sequence, using published sequence information (GenBank accession number X73874, incorporated herein as SEQ ID NO:3). For human Phosphorylase kinase alpha 1 the PCR  
15 primers were:

forward primer: GACTTCGGGATATGGGAACGT (SEQ ID NO: 4)

reverse primer: TCACACCAAACAGATCCAGTTCA (SEQ ID NO: 5) and

the PCR probe was: FAM-AACCAAGGGATCTCAGAGTTGAATGCCA-TAMRA

(SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City,  
20 CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:

forward primer: CAACGGATTGGTCGTATTGG (SEQ ID NO: 7)

reverse primer: GGCAACAATATCCACTTTACCAGAGT (SEQ ID NO: 8) and

25 the PCR probe was: 5' JOE-CGCCTGGTCACCAGGGCTGCT- TAMRA 3'

(SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

Probes and primers to mouse Phosphorylase kinase alpha 1  
30 were designed to hybridize to a mouse Phosphorylase kinase alpha 1 sequence, using published sequence information (GenBank accession number X74616, incorporated herein as SEQ ID NO:10). For mouse Phosphorylase kinase alpha 1 the PCR primers were:

-82-

forward primer: GGGAACGTGGCGATAAGACA (SEQ ID NO:11)  
reverse primer: ACCAAACAAGTCTAATTCATCTAGTGCTT (SEQ ID NO: 12)  
and the PCR probe was: FAM-CCAAGGCATCTCGGAATTGAATGCG-TAMRA  
(SEQ ID NO: 13) where FAM (PE-Applied Biosystems, Foster  
5 City, CA) is the fluorescent reporter dye) and TAMRA (PE-  
Applied Biosystems, Foster City, CA) is the quencher dye.  
For mouse GAPDH the PCR primers were:

forward primer: GGCAAATTCAACGGCACAGT (SEQ ID NO: 14)  
reverse primer: GGGTCTCGCTCCTGGAAGCT (SEQ ID NO: 15) and the  
10 PCR probe was: 5' JOE-AAGGCCGAGAATGGGAAGCTTGTCATC- TAMRA 3'  
(SEQ ID NO: 16) where JOE (PE-Applied Biosystems, Foster  
City, CA) is the fluorescent reporter dye) and TAMRA (PE-  
Applied Biosystems, Foster City, CA) is the quencher dye.

Probes and primers to rat Phosphorylase kinase alpha 1  
15 were designed to hybridize to a rat Phosphorylase kinase  
alpha 1 sequence, using published sequence information  
(GenBank accession number M92918, incorporated herein as SEQ  
ID NO:17). For rat Phosphorylase kinase alpha 1 the PCR  
primers were:

20 forward primer: TGAAGATGGAGCCAGCTTGA (SEQ ID NO: 18)  
reverse primer: CACCAAATAACCATCCTGCATT (SEQ ID NO: 19) and  
the PCR probe was: FAM-TCAAGTATCCTGGCAGCACTCCGGA-TAMRA  
(SEQ ID NO: 20) where FAM (PE-Applied Biosystems, Foster  
City, CA) is the fluorescent reporter dye) and TAMRA (PE-  
25 Applied Biosystems, Foster City, CA) is the quencher dye.

For rat0 GAPDH the PCR primers were:

forward primer: TGTTCTAGAGACAGCCGCATCTT (SEQ ID NO: 21)  
reverse primer: CACCGACCTTCACCATCTTGT (SEQ ID NO: 22) and the  
PCR probe was: 5' JOE-TTGTGCACTGCCAGCCTCGTCTCA- TAMRA 3' (SEQ  
30 ID NO: 23) where JOE (PE-Applied Biosystems, Foster City, CA)  
is the fluorescent reporter dye) and TAMRA (PE-Applied  
Biosystems, Foster City, CA) is the quencher dye.

**Example 14****Northern blot analysis of Phosphorylase kinase alpha 1 mRNA levels**

Eighteen hours after antisense treatment, cell  
5 monolayers were washed twice with cold PBS and lysed in 1 mL  
RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was  
prepared following manufacturer's recommended protocols.  
Twenty micrograms of total RNA was fractionated by  
electrophoresis through 1.2% agarose gels containing 1.1%  
10 formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon,  
OH). RNA was transferred from the gel to HYBOND™-N+ nylon  
membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by  
overnight capillary transfer using a Northern/Southern  
Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX).  
15 RNA transfer was confirmed by UV visualization. Membranes  
were fixed by UV cross-linking using a STRATALINKER™ UV  
Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then  
probed using QUICKHYB™ hybridization solution (Stratagene, La  
Jolla, CA) using manufacturer's recommendations for stringent  
20 conditions.

To detect human Phosphorylase kinase alpha 1, a human  
Phosphorylase kinase alpha 1 specific probe was prepared by  
PCR using the forward primer GACTTCGGGATATGGGAACGT (SEQ ID  
NO: 4) and the reverse primer TCACACCAAACAGATCCAGTTCA (SEQ ID  
25 NO: 5). To normalize for variations in loading and transfer  
efficiency membranes were stripped and probed for human  
glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA  
(Clontech, Palo Alto, CA).

To detect mouse Phosphorylase kinase alpha 1, a mouse  
30 Phosphorylase kinase alpha 1 specific probe was prepared by  
PCR using the forward primer GGGAACGTGGCGATAAGACA (SEQ ID  
NO:11) and the reverse primer ACCAAACAAGTCTAATTCATCTAGTGCTT  
(SEQ ID NO: 12). To normalize for variations in loading and  
transfer efficiency membranes were stripped and probed for

mouse glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

To detect rat Phosphorylase kinase alpha 1, a rat Phosphorylase kinase alpha 1 specific probe was prepared by  
5 PCR using the forward primer TGAAGATGGAGCCAGCTTGA (SEQ ID NO: 18) and the reverse primer CACCAAAATAACCATCCTGCATT (SEQ ID NO: 19). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for rat glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA  
10 (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### 15 Example 15

**Antisense inhibition of human Phosphorylase kinase alpha 1 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of  
20 oligonucleotides were designed to target different regions of the human Phosphorylase kinase alpha 1 RNA, using published sequences (GenBank accession number X73874, incorporated herein as SEQ ID NO: 3, and GenBank accession number X73878, incorporated herein as SEQ ID NO: 24). The oligonucleotides  
25 are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten  
30 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues

-85-

are 5-methylcytidines. The compounds were analyzed for their effect on human Phosphorylase kinase alpha 1 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present,  
 5 "N.D." indicates "no data".

Table 1

Inhibition of human Phosphorylase kinase alpha 1 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

10	ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%INHIB	SEQ ID NO
	118465	Start Codon	3	153	cggctcctcatggcgacacg	77	25
	118466	Coding	3	219	tgatggcacaggatggtctg	71	26
	118467	Coding	3	252	tagctggctggaagcaagcc	76	27
	118468	Coding	3	291	ctgtacacattatctcggac	90	28
15	118469	Coding	3	316	gcccacacccacacagcca	98	29
	118470	Coding	3	391	cttactacactctgctcca	89	30
	118471	Coding	3	426	tgtctgatcatgcagtgcag	93	31
	118472	Coding	3	438	actttatccacctgtctgat	91	32
	118473	Coding	3	549	tccaactgcagggtgtcccca	99	33
20	118474	Coding	3	600	agtcctgaggcagtcatttg	99	34
	118475	Coding	3	620	ctaggctgtggatgatatgg	86	35
	118476	Coding	3	675	gttttatatgcagcttcaat	50	36
	118477	Coding	3	811	cccacctttcacaccaaaca	90	37
	118478	Coding	3	834	aggacatggataactgattg	88	38
25	118479	Coding	3	1008	ccctgaagcttggtgatgat	85	39
	118480	Coding	3	1019	aaccataacgaccctgaagc	89	40
	118481	Coding	3	1056	ttaggagttttatatccatc	95	41
	118482	Coding	3	1104	aatagcttcagctcagctgg	81	42
	118483	Coding	3	1140	gtccagaacaatggccattc	98	43
30	118484	Coding	3	1196	tatattcttgaacctgttct	95	44
	118485	Coding	3	1230	ttgcccttgatgaggactgc	72	45
	118486	Coding	3	1350	ccccacatgtgaggcaattt	92	46
	118487	Coding	3	1360	tagagactgacccacatgt	94	47
	118488	Coding	3	1390	tccctctgccatcaagcttc	73	48
35	118489	Coding	3	1476	agaatggagacttgaaccac	85	49
	118490	Coding	3	1497	ttgatttcttctgtttcagc	96	50
	118491	Coding	3	1584	tggctgagaatacagagctgg	94	51
	118492	Coding	3	1631	gtccactgagtttcattcta	88	52
	118493	Coding	3	1704	gtgaaagtaaagatagtttt	39	53
40	118494	Coding	3	1714	aaactgtggagtgaagtaa	73	54
	118495	Coding	3	1734	tagaactgttgctggtctat	81	55
	118496	Coding	3	1746	tccagagccaggtagaactg	97	56
	118497	Coding	3	1803	cagcggctacagaggtagga	82	57
	118498	Coding	3	1825	ggtgggctggcctgtcatcc	91	58
45	118499	Coding	3	1868	ttccatcttcatcaagcatg	99	59
	118500	Coding	3	1890	aggatacttgaattcaagct	71	60
	118501	Coding	3	1909	catttttcggagtgtgcca	85	61
	118502	Coding	3	1959	tctgacaatttacctgtttg	99	62

- 86 -

	118503	Coding	3	1972	tggtgtcaaaaactctgaca	76	63
	118504	Coding	3	1977	caagatggtgtcaaaaactc	87	64
	118505	Coding	3	2010	gggccagggtccatgaagct	73	65
	118506	Coding	3	2021	gcttaccctcagggtccagg	96	66
5	118507	Coding	3	2041	atcataatcttcaactgtaca	76	67
	118508	Coding	3	2119	ttcatcaccacagcgagcat	80	68
	118509	Coding	3	2207	accgatctagccctcccttc	94	69
	118510	Coding	3	2250	accaaggacattaagtgcga	96	70
	118511	Coding	3	2293	atacatgtgaacattctgta	88	71
10	118512	Coding	3	2465	ttgagggtctccttcaactgt	94	72
	118513	Coding	3	2526	ttccagtcagggtcctttcat	91	73
	118514	Coding	3	2676	gtgcaggcctcatcaagtgc	30	74
	118515	Coding	3	2722	tggaggaagtccactgtca	97	75
	118516	Coding	3	2794	actggcttcacatcatcagct	95	76
15	118517	Coding	3	2837	ccattatttcctgtgtaagg	92	77
	118518	Coding	3	2926	tgcataacttgtatgatca	86	78
	118519	Coding	3	2973	tctgtggcttcctcagctga	57	79
	118520	Coding	3	2993	gactgagattcatcaggccc	93	80
	118521	Coding	3	3030	ctgagaatgtgatgcaggag	77	81
20	118522	Coding	3	3057	cttcgttccactccaaactc	63	82
	118523	Coding	3	3077	ttgaatcagtgggacgaacg	74	83
	118524	Coding	3	3123	gctccaacagcaccaatctc	94	84
	118525	Coding	3	3165	tcactttttaactgcatgat	80	85
	118526	Coding	3	3185	gaaattccacctgctttatc	17	86
25	118527	Coding	3	3295	ttgacgactatcttttagatg	67	87
	118528	Coding	3	3465	gagaattttaatctcacctgg	49	88
	118529	Coding	3	3489	ttcaggacagactccacatg	87	89
	118530	Coding	3	3540	acaaggatggcttcaaccag	93	90
	118531	Coding	3	3561	atatctgccagcatggtgag	86	91
30	118532	Coding	3	3571	atgaatttcaatatctgccca	92	92
	118533	Coding	3	3639	tggtcttgaaggacaagtc	67	93
	118534	Coding	3	3669	aacatgggtatcatctgcgcc	96	94
	118535	Coding	3	3679	atccttttgccaacatggtat	75	95
	118536	Coding	3	3702	agagtacagatgccagatgc	92	96
35	118537	Coding	3	3727	gccactgggtgcactgtcat	73	97
	118538	Coding	3	3750	aggtagggtcatggtgcaaaa	92	98
	118539	Coding	3	3783	aactcctgcacgtagggtggc	85	99
	118540	Stop	3	3823	ccaaagccctcattgcatgg	76	100
		Codon					
	118541	3'UTR	24	102	gtccaagctatgacaagcta	41	101
40	118542	3'UTR	24	351	ataatgggtattattaaaag	15	102

As shown in Table 1, SEQ ID NOs 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 76, 77, 78, 80, 81, 82, 83, 84, 85, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and 100 demonstrated at least 60% inhibition of human Phosphorylase kinase alpha 1 expression in this assay and are therefore preferred. The target sites to which these



preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

**Example 16**

**5 Antisense inhibition of mouse Phosphorylase kinase alpha 1 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap.**

In accordance with the present invention, a second series of oligonucleotides were designed to target different  
10 regions of the mouse Phosphorylase kinase alpha 1 RNA, using published sequences (GenBank accession number X74616, incorporated herein as SEQ ID NO: 10). The oligonucleotides are shown in Table 2. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to  
15 which the oligonucleotide binds. All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are  
20 composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on mouse Phosphorylase kinase alpha 1 mRNA levels by  
25 quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 2

Inhibition of mouse Phosphorylase kinase alpha 1 mRNA levels  
by chimeric phosphorothioate oligonucleotides having 2'-MOE  
wings and a deoxy gap

5

	ISIS #	REGION	TARGET	TARGET	SEQUENCE	%INHIB	SEQ ID
			SEQ ID NO	SITE			NO
10	118466	Coding	10	276	tgatggcacaggatggtctg	47	26
	118467	Coding	10	309	tagctggctggaagcaagcc	49	27
	118468	Coding	10	348	ctgtacacattatctcggac	63	28
	118469	Coding	10	373	gcccaaaccacacagcca	71	29
	118470	Coding	10	448	cttcactacactctgctcca	40	30
	118471	Coding	10	483	tgtctgatcatgcagtgcag	52	31
15	118472	Coding	10	495	actttatccacctgtctgat	69	32
	118473	Coding	10	606	tccaactgcagggtgtcccca	82	33
	118474	Coding	10	657	agtcctgaggcagtcatttg	62	34
	118475	Coding	10	677	ctaggctgtggatgatatgg	25	35
	118476	Coding	10	732	gttttatatgcagcttcaat	27	36
	118477	Coding	10	868	cccacctttcacaccaaaca	63	37
20	118478	Coding	10	891	aggacatggataaactgattg	52	38
	118479	Coding	10	1065	ccctgaagcttggtgatgat	66	39
	118480	Coding	10	1076	aaccataacgaccctgaagc	22	40
	118481	Coding	10	1113	ttaggagttttatatccatc	37	41
	118482	Coding	10	1161	aatagcttcagctcagctgg	52	42
	118483	Coding	10	1197	gtccagaacaatggccattc	73	43
25	118484	Coding	10	1253	tatattcttgaacctgttct	67	44
	118485	Coding	10	1287	ttgcccttgatgaggactgc	23	45
	118486	Coding	10	1407	ccccacatgtgaggcaattt	42	46
	118487	Coding	10	1417	tagagactgacccacatgt	29	47
	118488	Coding	10	1447	tccctctgccatcaagcttc	24	48
	118489	Coding	10	1533	agaatggagacttgaaccac	41	49
30	118490	Coding	10	1554	ttgatttcttctgtttcagc	43	50
	118491	Coding	10	1641	tggctgagaatacagagctgg	20	51
	118492	Coding	10	1688	gtccactgagtttcattcta	48	52
	118493	Coding	10	1761	gtgaaagtaaagatagtttt	0	53
	118494	Coding	10	1771	aaactgtggagtgaagtaa	42	54
	118495	Coding	10	1791	tagaactgttgctggtctat	46	55
35	118496	Coding	10	1803	tccagagccaggtagaactg	71	56
	118497	Coding	10	1860	cagcggctacagaggtagga	53	57
	118499	Coding	10	1925	ttccatcttcatcaagcatg	0	59
	118501	Coding	10	1966	catttttcggagtgtgcca	62	61
	118502	Coding	10	2016	tctgacaatttacctgtttg	33	62
	118503	Coding	10	2029	tgttgtcaaaaactctgaca	32	63
40	118504	Coding	10	2034	caagatgttgtcaaaaactc	0	64
	118505	Coding	10	2067	ggtccagggtccatgaagct	44	65
	118506	Coding	10	2078	gcttaccctcagggtccaggg	73	66
	118507	Coding	10	2098	atcataatcttcactgtaca	49	67
	118508	Coding	10	2179	ttcatcaccacagcgagcat	72	68
	118509	Coding	10	2267	accgatctagccctcccttc	64	69
45	118510	Coding	10	2310	accaaggacattaagtgcga	45	70
	118511	Coding	10	2353	atacatgtgaacattctgta	48	71
	118512	Coding	10	2525	ttgaggctctccttcaactgt	42	72
	118513	Coding	10	2586	ttccagtcagggtcctttcat	65	73

-89-

	118514	Coding	10	2736	gtgcaggcctcatcaagtgc	0	74
	118515	Coding	10	2782	tgagggaagtcctactgtca	42	75
	118516	Coding	10	2854	actggcttcacatcatcagct	69	76
	118517	Coding	10	2897	ccattatttcctgtgtaagg	41	77
5	118518	Coding	10	2986	tgccataacttgatgatca	52	78
	118519	Coding	10	3033	tctgtggcttcctcagctga	47	79
	118520	Coding	10	3053	gactgagattcatcaggccc	31	80
	118521	Coding	10	3090	ctgagaatgtgatgcaggag	57	81
	118522	Coding	10	3117	cttcgttcactccaaactc	4	82
10	118523	Coding	10	3137	ttgaatcagtgggacgaacg	18	83
	118524	Coding	10	3183	gctccaacagcaccaatctc	36	84
	118525	Coding	10	3225	tcactttttaactgcatgat	35	85
	118526	Coding	10	3245	gaaattccacctgctttatc	19	86
	118527	Coding	10	3406	ttgacgactatctttagatg	34	87
15	118528	Coding	10	3576	gagaattttaatctcacctgg	33	88
	118529	Coding	10	3600	ttcaggacagactccacatg	58	89
	118530	Coding	10	3651	acaaggatggcttcaaccag	53	90
	118531	Coding	10	3672	atatctgccagcatggtgag	54	91
	118532	Coding	10	3682	atgaatttcaatatctgcca	60	92
20	118533	Coding	10	3750	tggtcttgaaggaacaagtc	35	93
	118534	Coding	10	3780	aacatgggtatcatctgcgcc	60	94
	118535	Coding	10	3790	atcctttgccaacatgggtat	19	95
	118536	Coding	10	3813	agagtacagatgccagatgc	36	96
	118537	Coding	10	3838	gccactgggtgcactgtcat	36	97
25	118538	Coding	10	3861	aggtaggtcatggtgccaaa	59	98
	118539	Coding	10	3894	aactcctgcacgtaggtggc	46	99

As shown in Table 2, SEQ ID NOS 26, 27, 28, 29, 30, 31, 32, 33, 34, 37, 38, 39, 41, 42, 43, 44, 46, 49, 50, 52, 54, 55, 56, 57, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 76, 77, 78, 79, 80, 81, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 96, 97, 98 and 99 demonstrated at least 30% inhibition of mouse Phosphorylase kinase alpha 1 expression in this experiment and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

#### Example 17

Antisense inhibition of rat Phosphorylase kinase alpha 1 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap.

In accordance with the present invention, a third series of oligonucleotides were designed to target different regions

-90-

of the rat Phosphorylase kinase alpha 1 RNA, using published sequences (GenBank accession number M92918, incorporated herein as SEQ ID NO: 17). The oligonucleotides are shown in Table 3. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 3 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on rat Phosphorylase kinase alpha 1 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

20

Table 3

Inhibition of rat Phosphorylase kinase alpha 1 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

25	ISIS #	REGION	TARGET	TARGET	SEQUENCE	%INHIB	SEQ ID
			SEQ ID NO	SITE			NO
	118498	Coding	17	29	ggtgggctggcctgtcatcc	83	58
	118504	Coding	17	181	caagatgttgtcaaaaactc	73	64
	118545	Coding	17	5	cggctgcagaggtaggaga	88	103
	118546	Coding	17	9	tccaccggctgcagaggtag	85	104
30	118547	Coding	17	12	tcctccaccggctgcagagg	90	105
	118548	Coding	17	16	gtcatcctccaccggctgca	96	106
	118549	Coding	17	26	gggctggcctgtcatcctcc	91	107
	118550	Coding	17	35	agtgatggtgggctggcctg	84	108
	118551	Coding	17	37	aaagtgatggtgggctggcc	86	109
35	118552	Coding	17	58	agcatggtgtgtgagatggg	45	110
	118553	Coding	17	59	cagcatggtgtgtgagatgg	84	111
	118554	Coding	17	63	catccagcatggtgtgtgag	90	112
	118555	Coding	17	66	cttcatccagcatggtgtgt	94	113
	118556	Coding	17	67	tcttcatccagcatggtgtg	95	114

- 91 -

	118557	Coding	17	71	tccatcttcatccagcatgg	93	115
	118558	Coding	17	74	ggctccatcttcatccagca	93	116
	118559	Coding	17	77	gctggctccatcttcatcca	95	117
	118560	Coding	17	80	caagctggctccatcttcat	96	118
5	118561	Coding	17	87	ttgaattcaagctggctcca	89	119
	118562	Coding	17	93	ggatacttgaattcaagctg	88	120
	118563	Coding	17	95	caggatacttgaattcaagc	93	121
	118564	Coding	17	96	ccaggatacttgaattcaag	93	122
	118565	Coding	17	98	tgccaggatacttgaattca	93	123
10	118566	Coding	17	99	ctgccaggatacttgaattc	84	124
	118567	Coding	17	100	gctgccaggatacttgaatt	93	125
	118568	Coding	17	101	tgctgccaggatacttgaat	91	126
	118569	Coding	17	104	gagtgtgccaggatacttg	92	127
	118570	Coding	17	105	ggagtgtgccaggatactt	93	128
15	118571	Coding	17	108	tccggagtgtgccaggata	89	129
	118572	Coding	17	111	ttttccggagtgtgccagg	94	130
	118573	Coding	17	113	cattttccggagtgtgccca	91	131
	118574	Coding	17	114	gcattttccggagtgtgcc	94	132
	118575	Coding	17	117	cctgcattttccggagtgt	98	133
20	118576	Coding	17	120	catcctgcattttccggagt	98	134
	118577	Coding	17	123	aaccatcctgcattttccgg	99	135
	118578	Coding	17	126	aataaccatcctgcattttc	91	136
	118579	Coding	17	129	caaaataaaccatcctgcatt	81	137
	118580	Coding	17	131	accaaataaaccatcctgca	86	138
25	118581	Coding	17	133	ccaccaaataaaccatcctg	82	139
	118582	Coding	17	137	ggccccaccaaataaaccat	94	140
	118583	Coding	17	142	atcctggccccaccaaata	84	141
	118584	Coding	17	147	tttggatcctggccccacca	89	142
	118585	Coding	17	151	cctgtttggatcctggcccc	93	143
30	118586	Coding	17	154	ttacctgtttggatcctggc	88	144
	118587	Coding	17	157	agcttacctgtttggatcct	90	145
	118588	Coding	17	160	gacagcttacctgtttggat	88	146
	118589	Coding	17	163	tctgacagcttacctgtttg	87	147
	118590	Coding	17	166	aactctgacagcttacctgt	80	148
35	118591	Coding	17	169	aaaaactctgacagcttacc	91	149
	118592	Coding	17	172	gtcaaaaactctgacagctt	82	150
	118593	Coding	17	175	gttgtcaaaaactctgacag	87	151
	118594	Coding	17	178	gatgttgtcaaaaactctga	87	152
	118595	Coding	17	184	cagcaagatgttgtcaaaaa	89	153
40	118596	Coding	17	187	gtgcagcaagatgttgtcaa	89	154
	118597	Coding	17	190	tgtgtgcagcaagatgttgt	75	155
	118598	Coding	17	201	tgaagcttaagtgtgtgcag	84	156
	118599	Coding	17	208	gggtccatgaagcttaagt	73	157
	118600	Coding	17	212	tccagggtccatgaagctta	93	158
45	118601	Coding	17	241	taatcttcgctgtacagctt	78	159
	118602	Coding	17	251	gtcttcatcataatcttcgc	94	160
	118603	Coding	17	253	tagtcttcatcataatcttc	88	161
	118605	Coding	17	256	tcatagtcttcatcataatc	76	162
	118606	Coding	17	262	tcatcatcatagtcttcatc	90	163
50	118607	Coding	17	267	ccagctcatcatcatagtct	92	164
	118608	Coding	17	269	gtccagctcatcatcatagt	96	165
	118609	Coding	17	270	agtccagctcatcatcatag	95	166
	118610	Coding	17	272	agagtccagctcatcatcat	89	167
	118611	Coding	17	276	tgccagagtccagctcatca	94	168
55	118612	Coding	17	283	atccagttgccagagtccag	92	169
	118613	Coding	17	284	catccagttgccagagtcca	91	170

-92-

	118614	Coding	17	287	atccatccagttgccagagt	91	171
	118615	Coding	17	290	gctatccatccagttgccag	83	172
	118616	Coding	17	294	catagctatccatccagttg	89	173
	118617	Coding	17	298	gagtcatagctatccatcca	91	174
5	118618	Coding	17	304	cgtgttgagtcatagctatc	78	175
	118619	Coding	17	305	acgtgttgagtcatagctat	67	176
	118620	Coding	17	309	cattacgtgttgagtcatag	80	177
	118621	Coding	17	310	gcattacgtgttgagtcata	79	178

10 As shown in Table 3, SEQ ID NOs 58, 64, 103, 104, 105,  
 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118,  
 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130,  
 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142,  
 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154,  
 15 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166,  
 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177 and 178  
 demonstrated at least 60% inhibition of rat Phosphorylase  
 kinase alpha 1 expression in this experiment and are  
 therefore preferred. The target sites to which these  
 20 preferred sequences are complementary are herein referred to  
 as "active sites" and are therefore preferred sites for  
 targeting by compounds of the present invention.

#### Example 18

Western blot analysis of Phosphorylase kinase alpha 1 protein  
 25 levels

Western blot analysis (immunoblot analysis) is carried  
 out using standard methods. Cells are harvested 16-20 h  
 after oligonucleotide treatment, washed once with PBS,  
 suspended in Laemmli buffer (100 ul/well), boiled for 5  
 30 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for  
 1.5 hours at 150 V, and transferred to membrane for western  
 blotting. Appropriate primary antibody directed to  
 Phosphorylase kinase alpha 1 is used, with a radiolabelled or  
 fluorescently labeled secondary antibody directed against the  
 35 primary antibody species. Bands are visualized using a  
 PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

What is claimed is:

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding Phosphorylase kinase alpha 1, wherein said compound specifically hybridizes with and inhibits the expression of Phosphorylase kinase alpha 1.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 103, 104, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177 or 178.

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

- 94 -

10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding Phosphorylase kinase alpha 1.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of Phosphorylase kinase alpha 1 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of Phosphorylase kinase alpha 1 is inhibited.

16. A method of treating an animal having a disease or condition associated with Phosphorylase kinase alpha 1 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of Phosphorylase kinase alpha 1 is inhibited.

17. The method of claim 16 wherein the disease or condition is a metabolic disorder.

18. The method of claim 17 wherein the metabolic disorder is diabetes.



## SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.  
 Brett P. Monia  
 Jacqueline Wyatt

<120> ANTISENSE MODULATION OF PHOSPHORYLASE KINASE ALPHA 1 EXPRESSION

<130> RTSP-0165

<150> 09/657,452

<151> 2000-09-07

<160> 178

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctcttcaggg

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2

atgcattctg cccccaagga

20

<210> 3

<211> 4215

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (162)...(3833)

<400> 3

gccgccgggc gccaggcctg agcgggtggga gggctctgcg gggcctggtg ttcaggcgctc

60

ccaccacgag ggtggagcag cggtggatac ttgttcctta gggaccgaag ctccgggtggc

120

accggggcta tttctcagag gacaattagt aacgtgtcgc c atg agg agc cgg agt

176

Met Arg Ser Arg Ser  
 1 5

aac tcc ggg gtc cgg ctg gac ggc tac gct cga ctg gtg caa cag acc

224

Asn Ser Gly Val Arg Leu Asp Gly Tyr Ala Arg Leu Val Gln Gln Thr

10

15

20

atc ctg tgc cat cag aat cca gtg act ggc ttg ctt cca gcc agc tat

272

Ile Leu Cys His Gln Asn Pro Val Thr Gly Leu Leu Pro Ala Ser Tyr

25

30

35

gat cag aaa gat gct tgg gtc cga gat aat gtg tac agc atc ttg gct Asp Gln Lys Asp Ala Trp Val Arg Asp Asn Val Tyr Ser Ile Leu Ala 40 45 50	320
gtg tgg ggt ttg ggc ctg gcc tat cgg aag aat gca gac cgg gat gag Val Trp Gly Leu Gly Leu Ala Tyr Arg Lys Asn Ala Asp Arg Asp Glu 55 60 65	368
gat aag gca aag gcc tat gaa ttg gag cag agt gta gtg aag ctg atg Asp Lys Ala Lys Ala Tyr Glu Leu Glu Gln Ser Val Val Lys Leu Met 70 75 80 85	416
aga gga cta ctg cac tgc atg atc aga cag gtg gat aaa gta gaa tcc Arg Gly Leu Leu His Cys Met Ile Arg Gln Val Asp Lys Val Glu Ser 90 95 100	464
ttc aaa tat agt cag agt act aag gat agc ctc cat gca aag tac aac Phe Lys Tyr Ser Gln Ser Thr Lys Asp Ser Leu His Ala Lys Tyr Asn 105 110 115	512
acc aaa acc tgt gcc act gta gtg ggt gat gat caa tgg gga cac ctg Thr Lys Thr Cys Ala Thr Val Val Gly Asp Asp Gln Trp Gly His Leu 120 125 130	560
cag ttg gat gct acc tct gtg tac ctg ctc ttc tta gcc caa atg act Gln Leu Asp Ala Thr Ser Val Tyr Leu Leu Phe Leu Ala Gln Met Thr 135 140 145	608
gcc tca gga ctc cat atc atc cac agc cta gat gaa gtc aat ttc ata Ala Ser Gly Leu His Ile Ile His Ser Leu Asp Glu Val Asn Phe Ile 150 155 160 165	656
cag aac ctt gtg ttt tac att gaa gct gca tat aaa act gct gac ttc Gln Asn Leu Val Phe Tyr Ile Glu Ala Tyr Lys Thr Ala Asp Phe 170 175 180	704
ggg ata tgg gaa cgt gga gac aag acc aac caa ggg atc tca gag ttg Gly Ile Trp Glu Arg Gly Asp Lys Thr Asn Gln Gly Ile Ser Glu Leu 185 190 195	752
aat gcc agt tca gtt gga atg gca aag gca gcc ctg gaa gca tta gat Asn Ala Ser Ser Val Gly Met Ala Lys Ala Ala Leu Glu Ala Leu Asp 200 205 210	800
gaa ctg gat ctg ttt ggt gtg aaa ggt ggg cct caa tca gtt atc cat Glu Leu Asp Leu Phe Gly Val Lys Gly Gly Pro Gln Ser Val Ile His 215 220 225	848
gtc ctg gct gat gaa gta cag cac tgc cag tct atc cta aat tca cta Val Leu Ala Asp Glu Val Gln His Cys Gln Ser Ile Leu Asn Ser Leu 230 235 240 245	896
ctg ccc cgt gct tca aca tca aaa gag gtt gat gct agt cta ctc tca Leu Pro Arg Ala Ser Thr Ser Lys Glu Val Asp Ala Ser Leu Leu Ser 250 255 260	944
gtg gtt tcc ttc cct gcc ttt gca gta gag gat agc cag ttg gtg gag Val Val Ser Phe Pro Ala Phe Ala Val Glu Asp Ser Gln Leu Val Glu 265 270 275	992
ctc aca aaa cag gaa atc atc acc aag ctt cag ggt cgt tat ggt tgc Leu Thr Lys Gln Glu Ile Ile Thr Lys Leu Gln Gly Arg Tyr Gly Cys 280 285 290	1040
tgt cgc ttt cta cga gat gga tat aaa act cct aaa gag gat ccc aat	1088

Cys	Arg	Phe	Leu	Arg	Asp	Gly	Tyr	Lys	Thr	Pro	Lys	Glu	Asp	Pro	Asn		
295						300					305						
cg	ct	tac	tat	gaa	cca	gct	gag	ctg	aag	cta	ttt	gaa	aac	att	gag	1136	
Arg	Leu	Tyr	Tyr	Glu	Pro	Ala	Glu	Leu	Lys	Leu	Phe	Glu	Asn	Ile	Glu		
310				315						320					325		
tgt	gaa	tgg	cca	ttg	ttc	tgg	aca	tac	ttt	att	ctt	gat	ggg	gtc	ttc	1184	
Cys	Glu	Trp	Pro	Leu	Phe	Trp	Thr	Tyr	Phe	Ile	Leu	Asp	Gly	Val	Phe		
				330					335					340			
agt	ggc	aat	gca	gaa	cag	gtt	caa	gaa	tat	aaa	gag	gct	ctt	gaa	gca	1232	
Ser	Gly	Asn	Ala	Glu	Gln	Val	Gln	Glu	Tyr	Lys	Glu	Ala	Leu	Glu	Ala		
			345					350					355				
gtc	ctc	atc	aag	ggc	aaa	aat	gga	gtc	cca	ctt	ctg	cca	gag	ctg	tac	1280	
Val	Leu	Ile	Lys	Gly	Lys	Asn	Gly	Val	Pro	Leu	Leu	Pro	Glu	Leu	Tyr		
		360					365					370					
agt	gtt	cct	cct	gac	agg	gtc	gat	gaa	gaa	tat	cag	aat	cct	cac	act	1328	
Ser	Val	Pro	Pro	Asp	Arg	Val	Asp	Glu	Glu	Tyr	Gln	Asn	Pro	His	Thr		
	375					380					385						
gtg	gac	cga	gtc	ccc	atg	ggg	aaa	ttg	cct	cac	atg	tgg	ggg	cag	tct	1376	
Val	Asp	Arg	Val	Pro	Met	Gly	Lys	Leu	Pro	His	Met	Trp	Gly	Gln	Ser		
390					395					400					405		
cta	tac	att	tta	gga	agc	ttg	atg	gca	gag	gga	ttt	tta	gcc	cct	gga	1424	
Leu	Tyr	Ile	Leu	Gly	Ser	Leu	Met	Ala	Glu	Gly	Phe	Leu	Ala	Pro	Gly		
				410				415						420			
gaa	att	gat	ccc	ctg	aat	cgc	agg	ttt	tct	act	gta	ccg	aag	ccc	gat	1472	
Glu	Ile	Asp	Pro	Leu	Asn	Arg	Arg	Phe	Ser	Thr	Val	Pro	Lys	Pro	Asp		
			425					430					435				
gtt	gtg	gtt	caa	gtc	tcc	att	cta	gct	gaa	aca	gaa	gaa	atc	aag	acc	1520	
Val	Val	Val	Gln	Val	Ser	Ile	Leu	Ala	Glu	Thr	Glu	Glu	Ile	Lys	Thr		
		440					445					450					
att	ttg	aag	gac	aag	gga	att	tac	gtg	gag	acc	att	gct	gag	gta	tac	1568	
Ile	Leu	Lys	Asp	Lys	Gly	Ile	Tyr	Val	Glu	Thr	Ile	Ala	Glu	Val	Tyr		
	455					460					465						
ccc	atc	aga	gta	caa	cca	gct	cgt	att	ctc	agc	cac	att	tat	tcc	agc	1616	
Pro	Ile	Arg	Val	Gln	Pro	Ala	Arg	Ile	Leu	Ser	His	Ile	Tyr	Ser	Ser		
470					475					480					485		
cta	gga	tgc	aac	aat	aga	atg	aaa	ctc	agt	gga	cga	ccc	tac	aga	cac	1664	
Leu	Gly	Cys	Asn	Asn	Arg	Met	Lys	Leu	Ser	Gly	Arg	Pro	Tyr	Arg	His		
				490				495						500			
atg	gga	gtg	ctt	gga	act	tca	aaa	ctc	tat	gac	att	cgg	aaa	act	atc	1712	
Met	Gly	Val	Leu	Gly	Thr	Ser	Lys	Leu	Tyr	Asp	Ile	Arg	Lys	Thr	Ile		
			505					510					515				
ttt	act	ttc	act	cca	cag	ttt	ata	gac	cag	caa	cag	ttc	tac	ctg	gct	1760	
Phe	Thr	Phe	Thr	Pro	Gln	Phe	Ile	Asp	Gln	Gln	Gln	Phe	Tyr	Leu	Ala		
		520					525					530					
ctg	gac	aac	aag	atg	ata	gtg	gaa	atg	ctt	aga	aca	gac	ctc	tcc	tac	1808	
Leu	Asp	Asn	Lys	Met	Ile	Val	Glu	Met	Leu	Arg	Thr	Asp	Leu	Ser	Tyr		
	535					540					545						
ctc	tgt	agc	cgc	tgg	cgg	atg	aca	ggc	cag	ccc	acc	atc	acc	ttc	ccc	1856	
Leu	Cys	Ser	Arg	Trp	Arg	Met	Thr	Gly	Gln	Pro	Thr	Ile	Thr	Phe	Pro		

550	555	560	565	
atc tca tac agc atg ctt gat gaa gat gga aca agc ttg aat tca agt Ile Ser Tyr Ser Met Leu Asp Glu Asp Gly Thr Ser Leu Asn Ser Ser 570 575 580				1904
atc ctg gca gca ctc cga aaa atg caa gat ggg tat ttt ggt ggg gca Ile Leu Ala Ala Leu Arg Lys Met Gln Asp Gly Tyr Phe Gly Gly Ala 585 590 595				1952
agg gtt caa aca ggt aaa ttg tca gag ttt ttg aca aca tct tgt tgc Arg Val Gln Thr Gly Lys Leu Ser Glu Phe Leu Thr Thr Ser Cys Cys 600 605 610				2000
aca cac ttg agc ttc atg gac cct gga cct gag ggt aag ctg tac agt Thr His Leu Ser Phe Met Asp Pro Gly Pro Glu Gly Lys Leu Tyr Ser 615 620 625				2048
gaa gat tat gat gac aac tat gat tac ctg gaa tct ggc aac tgg atg Glu Asp Tyr Asp Asp Asn Tyr Asp Tyr Leu Glu Ser Gly Asn Trp Met 630 635 640 645				2096
aat gat tat gat tca acc agt cat gct cgc tgt ggt gat gaa gtt gct Asn Asp Tyr Asp Ser Thr Ser His Ala Arg Cys Gly Asp Glu Val Ala 650 655 660				2144
cgt tat tta gat cac ctt ttg gcg cac act gct ccc cat cct aaa cta Arg Tyr Leu Asp His Leu Leu Ala His Thr Ala Pro His Pro Lys Leu 665 670 675				2192
gcc cct acc tca cag aag gga ggg cta gat cgg ttc caa gct gct gtg Ala Pro Thr Ser Gln Lys Gly Gly Leu Asp Arg Phe Gln Ala Ala Val 680 685 690				2240
caa aca acc tgc gac tta atg tcc ttg gtg acc aag gcc aag gaa ctg Gln Thr Thr Cys Asp Leu Met Ser Leu Val Thr Lys Ala Lys Glu Leu 695 700 705				2288
cat gta cag aat gtt cac atg tat ctt cct acg aag tta ttt cag gct His Val Gln Asn Val His Met Tyr Leu Pro Thr Lys Leu Phe Gln Ala 710 715 720 725				2336
tcc cgg cct tca ttc aac tta ctt gat tca cct cat ccc cga cag gag Ser Arg Pro Ser Phe Asn Leu Leu Asp Ser Pro His Pro Arg Gln Glu 730 735 740				2384
aac cag gtt ccc tct gtt cgt gta gaa ata cat ctt cct aga gac cag Asn Gln Val Pro Ser Val Arg Val Glu Ile His Leu Pro Arg Asp Gln 745 750 755				2432
tct ggg gag gtg gac ttt aaa gca ctg gtt tta cag ttg aag gag acc Ser Gly Glu Val Asp Phe Lys Ala Leu Val Leu Gln Leu Lys Glu Thr 760 765 770				2480
tca agc tta cag gaa caa gct gat atc ctc tat atg ctg tat act atg Ser Ser Leu Gln Glu Gln Ala Asp Ile Leu Tyr Met Leu Tyr Thr Met 775 780 785				2528
aaa gga cct gac tgg aac act gaa ttg tat aat gaa cgg agt gct aca Lys Gly Pro Asp Trp Asn Thr Glu Leu Tyr Asn Glu Arg Ser Ala Thr 790 795 800 805				2576
gtg aga gag ctt ctt acc gag ctg tat ggc aaa gtg gga gaa att cgt Val Arg Glu Leu Leu Thr Glu Leu Tyr Gly Lys Val Gly Glu Ile Arg 810 815 820				2624

cac tgg ggc ctg atc cga tac att tct ggg atc tta agg aag aaa gtg His Trp Gly Leu Ile Arg Tyr Ile Ser Gly Ile Leu Arg Lys Lys Val 825 830 835	2672
gaa gca ctt gat gag gcc tgc aca gac ctt ctc tcc cac cag aaa cat Glu Ala Leu Asp Glu Ala Cys Thr Asp Leu Leu Ser His Gln Lys His 840 845 850	2720
ttg aca gta gga ctt cct cca gaa cct cga gaa aag act atc tct gca Leu Thr Val Gly Leu Pro Pro Glu Pro Arg Glu Lys Thr Ile Ser Ala 855 860 865	2768
cct ctg ccc tat gag gcg ctc act cag ctg ata gat gaa gcc agt gaa Pro Leu Pro Tyr Glu Ala Leu Thr Gln Leu Ile Asp Glu Ala Ser Glu 870 875 880 885	2816
ggg gat atg agc att tca atc ctt aca cag gaa ata atg gta tat cta Gly Asp Met Ser Ile Ser Ile Leu Thr Gln Glu Ile Met Val Tyr Leu 890 895 900	2864
gcc atg tat atg cga acc cag cct ggc ctc ttt gct gaa atg ttt cga Ala Met Tyr Met Arg Thr Gln Pro Gly Leu Phe Ala Glu Met Phe Arg 905 910 915	2912
ctt cga att ggt ctg atc ata caa gtt atg gca aca gaa ctg gcc cac Leu Arg Ile Gly Leu Ile Ile Gln Val Met Ala Thr Glu Leu Ala His 920 925 930	2960
tcc ctt cga tgc tca gct gag gaa gcc aca gag ggc ctg atg aat ctc Ser Leu Arg Cys Ser Ala Glu Glu Ala Thr Glu Gly Leu Met Asn Leu 935 940 945	3008
agt cct tcg gcc atg aag aat ctc ctg cat cac att ctc agc ggc aag Ser Pro Ser Ala Met Lys Asn Leu Leu His His Ile Leu Ser Gly Lys 950 955 960 965	3056
gag ttt gga gtg gaa cga agc gtt cgt ccc act gat tca aat gtc agt Glu Phe Gly Val Glu Arg Ser Val Arg Pro Thr Asp Ser Asn Val Ser 970 975 980	3104
cct gct att tct atc cac gag att ggt gct gtt gga gca acc aaa aca Pro Ala Ile Ser Ile His Glu Ile Gly Ala Val Gly Ala Thr Lys Thr 985 990 995	3152
gaa cga act ggg atc atg cag tta aaa agt gag ata aag cag gtg gaa Glu Arg Thr Gly Ile Met Gln Leu Lys Ser Glu Ile Lys Gln Val Glu 1000 1005 1010	3200
ttt cgt aga ctg tca atc tca gct gag agt cag tca cct gga acc tct Phe Arg Arg Leu Ser Ile Ser Ala Glu Ser Gln Ser Pro Gly Thr Ser 1015 1020 1025	3248
atg act cca agt agt ggg tcc ttt cct agt gca tat gat cag cag tca Met Thr Pro Ser Ser Gly Ser Phe Pro Ser Ala Tyr Asp Gln Gln Ser 1030 1035 1040 1045	3296
tct aaa gat agt cgt caa ggt caa tgg caa cgc cga aga agg ctg gat Ser Lys Asp Ser Arg Gln Gly Gln Trp Gln Arg Arg Arg Arg Leu Asp 1050 1055 1060	3344
ggg gca ctg aat aga gtt cca gtt gga ttt tat cag aaa gta tgg aaa Gly Ala Leu Asn Arg Val Pro Val Gly Phe Tyr Gln Lys Val Trp Lys 1065 1070 1075	3392
gtt ttg cag aag tgt cac gga ctt tct gtt gaa ggg ttt gtc ctt cct	3440

Val Leu Gln Lys Cys His Gly Leu Ser Val Glu Gly Phe Val Leu Pro.  
1080 1085 1090

tcc tct acc act aga gag atg act cca ggt gag att aaa ttc tct gtt 3488  
Ser Ser Thr Thr Arg Glu Met Thr Pro Gly Glu Ile Lys Phe Ser Val  
1095 1100 1105

cat gtg gag tct gtc ctg aat cgt gta cct cag cca gag tac cgt cag 3536  
His Val Glu Ser Val Leu Asn Arg Val Pro Gln Pro Glu Tyr Arg Gln  
1110 1115 1120 1125

ctg ctg gtt gaa gcc atc ctt gtc ctc acc atg ctg gca gat att gaa 3584  
Leu Leu Val Glu Ala Ile Leu Val Leu Thr Met Leu Ala Asp Ile Glu  
1130 1135 1140

att cat agc atc gga agc atc att gct gtg gaa aaa ata gtg cat att 3632  
Ile His Ser Ile Gly Ser Ile Ile Ala Val Glu Lys Ile Val His Ile  
1145 1150 1155

gcc aat gac ttg ttc ctt caa gaa cag aaa acc ctt ggc gca gat gat 3680  
Ala Asn Asp Leu Phe Leu Gln Glu Gln Lys Thr Leu Gly Ala Asp Asp  
1160 1165 1170

acc atg ttg gca aag gat ccc gca tct ggc atc tgt act ctt ctg tat 3728  
Thr Met Leu Ala Lys Asp Pro Ala Ser Gly Ile Cys Thr Leu Leu Tyr  
1175 1180 1185

gac agt gca ccc agt ggc agg ttt ggc acc atg acc tac ctc tcc aag 3776  
Asp Ser Ala Pro Ser Gly Arg Phe Gly Thr Met Thr Tyr Leu Ser Lys  
1190 1195 1200 1205

gca gcc gcc acc tac gtg cag gag ttc ctg ccc cac agc atc tgt gcc 3824  
Ala Ala Ala Thr Tyr Val Gln Glu Phe Leu Pro His Ser Ile Cys Ala  
1210 1215 1220

atg caa tga gggctttggt tcctggcttc tgggagcctt ttgacagctg gtccctgcct 3883  
Met Gln

cggttgattg tgcatggaac taaaatgtta ttgcctaata actccaaccc tgcccccttc 3943

tgtcccatcc ttcccaagaa gagagaactt ttctgataaa ctaactactg tagaagaagt 4003

gaacacttac ctggaggctc accttcgaga accagtgaca atcttatgag tataatgaac 4063

actcagccag gcctgtcatg attggcttta tttctttcat cattcataaa agtttgcatg 4123

tgtttttatt ctctagatct gttaccaata tagttttcta actcctgttt ggggagcaag 4183

tgtaataat aacttattcc taaaaaaaaa aa 4215

<210> 4  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 4  
gacttcggga tatgggaacg t 21

<210> 5

<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 5  
tcacaccaaa cagatccagt tca 23

<210> 6  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 6  
aaccaaggga tctcagagtt gaatgcca 28

<210> 7  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 7  
caacggattt ggtcgtattg g 21

<210> 8  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 8  
ggcaacaata tccactttac cagagt 26

<210> 9  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 9  
cgcctgggtca ccagggtgc t 21

<210> 10  
<211> 4130  
<212> DNA  
<213> Mus musculus

<220>

&lt;221&gt; CDS

&lt;222&gt; (219)...(3944)

&lt;400&gt; 10

```

cggggaccac gggagactga gtggagacgg ggtctttgcc ggccgaagcc tagaccgcgg      60
gaggctgagc cgcggaatga agtggactct gaggccgccg gaaccagggtg acgcgcggga      120
accattggcc acacgagcaa gggagcgttt agaaggtttc cgtaggcctg gcgtccgagc      180
ttcccgtgtg tgcagtgggg gattagggac gtggcacc atg agg agc cgc agt aac      236
                               Met Arg Ser Arg Ser Asn
                               1           5

tcg gga gtc cgc ctg gac ggc tat gct cgg ctg gtg cat cag acc atc      284
Ser Gly Val Arg Leu Asp Gly Tyr Ala Arg Leu Val His Gln Thr Ile
                               10           15           20

ctg tgc cat cag aat cca gtg aca ggc ttg ctt cca gcc agc tat gat      332
Leu Cys His Gln Asn Pro Val Thr Gly Leu Leu Pro Ala Ser Tyr Asp
                               25           30           35

caa aaa gat gcc tgg gtc cga gat aat gtg tac agt atc ctg gct gtg      380
Gln Lys Asp Ala Trp Val Arg Asp Asn Val Tyr Ser Ile Leu Ala Val
                               40           45           50

tgg ggt ttg ggc ctg gca tat cgc aaa aat gct gac cgt gat gag gac      428
Trp Gly Leu Gly Leu Ala Tyr Arg Lys Asn Ala Asp Arg Asp Glu Asp
                               55           60           65           70

aag gca aag gcc tat gaa ctg gag cag agt gta gtg aag tta atg agg      476
Lys Ala Lys Ala Tyr Glu Leu Glu Gln Ser Val Val Lys Leu Met Arg
                               75           80           85

gga ctg ctg cac tgc atg atc aga cag gtg gat aaa gta gag tcc ttc      524
Gly Leu Leu His Cys Met Ile Arg Gln Val Asp Lys Val Glu Ser Phe
                               90           95           100

aag tac agt cag agt act ccc gat agc ctc cat gcc aag tac aac acc      572
Lys Tyr Ser Gln Ser Thr Pro Asp Ser Leu His Ala Lys Tyr Asn Thr
                               105           110           115

aaa act tgt gcc act gtg gtg ggt gat gac cag tgg gga cac ctg cag      620
Lys Thr Cys Ala Thr Val Val Gly Asp Asp Gln Trp Gly His Leu Gln
                               120           125           130

ttg gat gct act tct gtg tac ctg ctc ttc cta gca caa atg act gcc      668
Leu Asp Ala Thr Ser Val Tyr Leu Leu Phe Leu Ala Gln Met Thr Ala
                               135           140           145           150

tca gga ctc cat atc atc cac agc cta gac gaa gtc aat ttt ata cag      716
Ser Gly Leu His Ile Ile His Ser Leu Asp Glu Val Asn Phe Ile Gln
                               155           160           165

aac ctc gtg ttt tac att gaa gct gca tat aaa act gct gac ttt ggg      764
Asn Leu Val Phe Tyr Ile Glu Ala Ala Tyr Lys Thr Ala Asp Phe Gly
                               170           175           180

ata tgg gaa cgt ggc gat aag aca aac caa ggc atc tcg gaa ttg aat      812
Ile Trp Glu Arg Gly Asp Lys Thr Asn Gln Gly Ile Ser Glu Leu Asn
                               185           190           195

gcg agt tca gtt gga atg gcc aag gca gcc ctg gaa gca cta gat gaa      860
Ala Ser Ser Val Gly Met Ala Lys Ala Ala Leu Glu Ala Leu Asp Glu
                               200           205           210

```



tta gac ttg ttt ggt gtg aaa ggt ggg cca caa tca gtt atc cat gtc	908
Leu Asp Leu Phe Gly Val Lys Gly Gly Pro Gln Ser Val Ile His Val	
215 220 225 230	
ctg gct gat gaa gtc caa cac tgc cag tct atc ctg aat tca cta ctg	956
Leu Ala Asp Glu Val Gln His Cys Gln Ser Ile Leu Asn Ser Leu Leu	
235 240 245	
ccc agg gct tca aca tcc aaa gaa gtt gat gcc agt ctg ctc tca gtg	1004
Pro Arg Ala Ser Thr Ser Lys Glu Val Asp Ala Ser Leu Leu Ser Val	
250 255 260	
gtc tct ttc cca gcc ttt gct gta gag gac agc cat ttg gtg gag ctc	1052
Val Ser Phe Pro Ala Phe Ala Val Glu Asp Ser His Leu Val Glu Leu	
265 270 275	
acc aaa cag gag atc atc acc aag ctt cag ggt cgt tat ggt tgc tgt	1100
Thr Lys Gln Glu Ile Ile Thr Lys Leu Gln Gly Arg Tyr Gly Cys Cys	
280 285 290	
cgt ttt ctg cga gat gga tat aaa act cct aaa gag gat ccc caa cgc	1148
Arg Phe Leu Arg Asp Gly Tyr Lys Thr Pro Lys Glu Asp Pro Gln Arg	
295 300 305 310	
cta tac tat aac cca gct gag ctg aag cta ttt gaa aac att gag tgc	1196
Leu Tyr Tyr Asn Pro Ala Glu Leu Lys Leu Phe Glu Asn Ile Glu Cys	
315 320 325	
gaa tgg cca ttg ttc tgg aca tac ttt atc ctt gat ggg atc ttc agt	1244
Glu Trp Pro Leu Phe Trp Thr Tyr Phe Ile Leu Asp Gly Ile Phe Ser	
330 335 340	
ggc aac gta gaa cag gtt caa gaa tat aga gag gct ctt gat gca gtc	1292
Gly Asn Val Glu Gln Val Gln Glu Tyr Arg Glu Ala Leu Asp Ala Val	
345 350 355	
ctc atc aag ggc aaa aat gga gtc cct ctt ctt cca gag ctg tac agt	1340
Leu Ile Lys Gly Lys Asn Gly Val Pro Leu Leu Pro Glu Leu Tyr Ser	
360 365 370	
gtc cct cct gac agg gtt gat gaa gag tat caa aat ccc cac act gtg	1388
Val Pro Pro Asp Arg Val Asp Glu Glu Tyr Gln Asn Pro His Thr Val	
375 380 385 390	
gat cga gtc cct atg gga aaa ttg cct cac atg tgg ggt cag tct cta	1436
Asp Arg Val Pro Met Gly Lys Leu Pro His Met Trp Gly Gln Ser Leu	
395 400 405	
tac att tta gga agc ttg atg gca gag gga ttt tta gct cct gga gaa	1484
Tyr Ile Leu Gly Ser Leu Met Ala Glu Gly Phe Leu Ala Pro Gly Glu	
410 415 420	
att gat ccc ctg aat cgt agg ttt tct act gtg cca aag cca gat gtg	1532
Ile Asp Pro Leu Asn Arg Arg Phe Ser Thr Val Pro Lys Pro Asp Val	
425 430 435	
gtg gtt caa gtc tcc att ctg gct gaa aca gaa gaa atc aag gcc att	1580
Val Val Gln Val Ser Ile Leu Ala Glu Thr Glu Glu Ile Lys Ala Ile	
440 445 450	
ttg aag gac aaa gga att gat gtg gag acc att gct gaa gtg tac ccc	1628
Leu Lys Asp Lys Gly Ile Asp Val Glu Thr Ile Ala Glu Val Tyr Pro	
455 460 465 470	
ata aga gta cag cca gct cgt att ctc agc cat att tat tct agt cta	1676

Ile Arg Val Gln Pro Ala Arg Ile Leu Ser His Ile Tyr Ser Ser Leu	
475 480 485	
gga tgc aac agt aga atg aaa ctc agt gga cga ccc tac agg ctc atg	1724
Gly Cys Asn Ser Arg Met Lys Leu Ser Gly Arg Pro Tyr Arg Leu Met	
490 495 500	
ggg gtg ctt gga aca tca aaa ctt tat gac att cgc aaa act atc ttt	1772
Gly Val Leu Gly Thr Ser Lys Leu Tyr Asp Ile Arg Lys Thr Ile Phe	
505 510 515	
act ttc act cca cag ttt ata gac cag caa cag ttc tac ctg gct ctg	1820
Thr Phe Thr Pro Gln Phe Ile Asp Gln Gln Gln Phe Tyr Leu Ala Leu	
520 525 530	
gac aac cag atg ata gta gaa atg ctc aga aca gac ctt tcc tac ctc	1868
Asp Asn Gln Met Ile Val Glu Met Leu Arg Thr Asp Leu Ser Tyr Leu	
535 540 545 550	
tgt agc cgc tgg agg atg aca ggc cag ccc acg atc act ttc cct atc	1916
Cys Ser Arg Trp Arg Met Thr Gly Gln Pro Thr Ile Thr Phe Pro Ile	
555 560 565	
tgc cac acc atg ctt gat gaa gat gga acc agc ttg aat tca agt atc	1964
Ser His Thr Met Leu Asp Glu Asp Gly Thr Ser Leu Asn Ser Ser Ile	
570 575 580	
ttg gca gca ctc cga aaa atg cag gat ggc tat ttt ggt ggg gcc agg	2012
Leu Ala Ala Leu Arg Lys Met Gln Asp Gly Tyr Phe Gly Gly Ala Arg	
585 590 595	
atc caa aca ggt aaa ttg tca gag ttt ttg aca aca tct tgc tgc aca	2060
Ile Gln Thr Gly Lys Leu Ser Glu Phe Leu Thr Thr Ser Cys Cys Thr	
600 605 610	
cac tta agc ttc atg gac cct gga cct gag ggt aag ctg tac agt gaa	2108
His Leu Ser Phe Met Asp Pro Gly Pro Glu Gly Lys Leu Tyr Ser Glu	
615 620 625 630	
gat tat gat gaa gac tat gaa gat gat ttg gac tct ggc aac tgg atg	2156
Asp Tyr Asp Glu Asp Tyr Glu Asp Asp Leu Asp Ser Gly Asn Trp Met	
635 640 645	
gac agc tat gat tca aca agt aat gct cgc tgt ggt gat gaa gtt gcc	2204
Asp Ser Tyr Asp Ser Thr Ser Asn Ala Arg Cys Gly Asp Glu Val Ala	
650 655 660	
cgt tat tta gac cgc ctt ttg gca cac act gtt ccc cat cct aaa cta	2252
Arg Tyr Leu Asp Arg Leu Leu Ala His Thr Val Pro His Pro Lys Leu	
665 670 675	
gct cct acc tca cgg aag gga ggg cta gat cgg ttc cga gct gct gtg	2300
Ala Pro Thr Ser Arg Lys Gly Gly Leu Asp Arg Phe Arg Ala Ala Val	
680 685 690	
caa aca act tgc gac tta atg tcc ttg gtg gcc aag gcc aag gag ctg	2348
Gln Thr Thr Cys Asp Leu Met Ser Leu Val Ala Lys Ala Lys Glu Leu	
695 700 705 710	
cat ata cag aat gtt cac atg tat att cct aca aag tta ttt cag cct	2396
His Ile Gln Asn Val His Met Tyr Ile Pro Thr Lys Leu Phe Gln Pro	
715 720 725	
tct cgt cct tca ctc aac tta ctt gac toc cct gag tct cca caa gat	2444
Ser Arg Pro Ser Leu Asn Leu Leu Asp Ser Pro Glu Ser Pro Gln Asp	

11/44

gaa cga act gga atc atg cag tta aaa agt gag ata aag cag gtg gaa 3260  
 Glu Arg Thr Gly Ile Met Gln Leu Lys Ser Glu Ile Lys Gln Val Glu  
 1000 1005 1010

ttt cgt agg ctg tct gtc tcg atg gag agt cag act agt ggt ggt cat 3308  
 Phe Arg Arg Leu Ser Val Ser Met Glu Ser Gln Thr Ser Gly Gly His  
 1015 1020 1025 1030

ccc tcg ggt gta gat ttg atg tcg cca tcc ttt ctg tcc cct gca gcc 3356  
 Pro Ser Gly Val Asp Leu Met Ser Pro Ser Phe Leu Ser Pro Ala Ala  
 1035 1040 1045

tgt att gct gca agt agt gga tcc ttt cct acg gtg tgt gat cat cag 3404  
 Cys Ile Ala Ser Ser Gly Ser Phe Pro Thr Val Cys Asp His Gln  
 1050 1055 1060

aca tct aaa gat agt cgt caa ggc cag tgg caa cgc agg aga agg cta 3452  
 Thr Ser Lys Asp Ser Arg Gln Gln Trp Gln Arg Arg Arg Leu  
 1065 1070 1075

gat gga gca cta aat aga gta cca att gga ttt tat caa aaa gta tgg 3500  
 Asp Gly Ala Leu Asn Arg Val Pro Ile Gly Phe Tyr Gln Lys Val Trp  
 1080 1085 1090

aaa att ttg cag aaa tgt cat ggg ctt tct gtg gaa ggt ttt gtt ctt 3548  
 Lys Ile Leu Gln Lys Cys His Gly Leu Ser Val Glu Gly Phe Val Leu  
 1095 1100 1105 1110

ccc tct tca acc act agg gag atg acc cca ggt gag att aaa ttc tct 3596  
 Pro Ser Ser Thr Thr Arg Glu Met Thr Pro Gly Glu Ile Lys Phe Ser  
 1115 1120 1125

gtc cat gtg gag tct gtc ctg aat cgt gtc cct cag cca gaa tac cgc 3644  
 Val His Val Glu Ser Val Leu Asn Arg Val Pro Gln Pro Glu Tyr Arg  
 1130 1135 1140

caa ctt ctg gtt gaa gcc atc ctt gtt ctc acc atg ctg gca gat att 3692  
 Gln Leu Leu Val Glu Ala Ile Leu Val Leu Thr Met Leu Ala Asp Ile  
 1145 1150 1155

gaa att cat agc att ggg agc atc att gct gtg gag aaa ata gtt cat 3740  
 Glu Ile His Ser Ile Gly Ser Ile Ile Ala Val Glu Lys Ile Val His  
 1160 1165 1170

att gcc aac gac ttg ttc ctt caa gaa cag aaa acc ctc ggc gca gat 3788  
 Ile Ala Asn Asp Leu Phe Leu Gln Glu Gln Lys Thr Leu Gly Ala Asp  
 1175 1180 1185 1190

gat acc atg ttg gca aag gat cct gca tct ggc atc tgt act ctc ctg 3836  
 Asp Thr Met Leu Ala Lys Asp Pro Ala Ser Gly Ile Cys Thr Leu Leu  
 1195 1200 1205

tat gac agt gca ccc agt ggc aga ttt ggc acc atg acc tac ctc tcc 3884  
 Tyr Asp Ser Ala Pro Ser Gly Arg Phe Gly Thr Met Thr Tyr Leu Ser  
 1210 1215 1220

aag gca gct gcc acc tac gtg cag gag ttt ctg cca cac agc ctc tgt 3932  
 Lys Ala Ala Ala Thr Tyr Val Gln Glu Phe Leu Pro His Ser Leu Cys  
 1225 1230 1235

gcc atg cag tga gggctgtggg ttctgcctgt tagaacactt cagtagctag 3984  
 Ala Met Gln  
 1240

cccgtctggg tgattgtgca tggaactgac atgttaacac ctgatcactc cccaggccc 4044

ttttcctgcc ccatccctcc aggaagagag aactttgctg acgatctcac tacttttgaa 4104  
gaaaagaaga aatatctgga gacttg 4130

<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 11  
gggaacgtgg cgataagaca 20

<210> 12  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 12  
accaaacaag totaattcat ctagtgctt 29

<210> 13  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 13  
ccaaggcatc tcggaattga atgcg 25

<210> 14  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 14  
ggcaaattca acggcacagt 20

<210> 15  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 15  
gggtctcgct cctggaagct 20

<210> 16  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Probe

<400> 16  
 aagggcgcgaga atggggaagct tgtcctc

27

<210> 17  
 <211> 336  
 <212> DNA  
 <213> Rattus norvegicus

<220>  
 <221> CDS  
 <222> (1) ... (336)

<400> 17  
 gat ctc tcc tac ctc tgc agc cgg tgg agg atg aca ggc cag ccc acc  
 48  
 Asp Leu Ser Tyr Leu Cys Ser Arg Trp Arg Met Thr Gly Gln Pro Thr  
 1 5 10 15

atc act ttc ccc atc tca cac acc atg ctg gat gaa gat gga gcc agc  
 Ile Thr Phe Pro Ile Ser His Thr Met Leu Asp Glu Asp Gly Ala Ser  
 20 25 30

96

ttg aat tca agt atc ctg gca gca ctc cgg aaa atg cag gat ggt tat  
 Leu Asn Ser Ser Ile Leu Ala Ala Leu Arg Lys Met Gln Asp Gly Tyr  
 35 40 45

144

ttt ggt ggg gcc agg atc caa aca ggt aag ctg tca gag ttt ttg aca  
 Phe Gly Gly Ala Arg Ile Gln Thr Gly Lys Leu Ser Glu Phe Leu Thr  
 50 55 60

192

aca tct tgc tgc aca cac tta agc ttc atg gac cct gga ccc gag ggt  
 Thr Ser Cys Cys Thr His Leu Ser Phe Met Asp Pro Gly Pro Glu Gly  
 65 70 75 80

240

aag ctg tac agc gaa gat tat gat gaa gac tat gat gat gag ctg gac  
 Lys Leu Tyr Ser Glu Asp Tyr Asp Glu Asp Tyr Asp Asp Glu Leu Asp  
 85 90 95

288

tct ggc aac tgg atg gat agc tat gac tca aca cgt aat gct cgc tgt  
 Ser Gly Asn Trp Met Asp Ser Tyr Asp Ser Thr Arg Asn Ala Arg Cys  
 100 105 110

336

336

<210> 18  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> PCR Primer

<400> 18  
 tgaagatgga gccagcttga

20

<210> 19  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 19  
caccaaaaata accatcctgc att

23

<210> 20  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 20  
tcaagtatcc tggcagcact ccgga

25

<210> 21  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 21  
tgtttctagag acagccgcat ctt

23

<210> 22  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Primer

<400> 22  
caccgacctt caccatcttg t

21

<210> 23  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> PCR Probe

<400> 23  
ttgtgcagtg ccagcctcgt ctca

24

<210> 24  
<211> 417  
<212> DNA  
<213> Homo sapiens

<220>  
 <221> CDS  
 <222> (1)...(60)

<400> 24  
 tcc ggg gtc cgg ctg gac ggc tac gct cga ctg gtg caa cag acc atc  
 48  
 Ser Gly Val Arg Leu Asp Gly Tyr Ala Arg Leu Val Gln Gln Thr Ile  
 1 5 10 15  
 ctg tgc cat cag gtaactgcag ggggtgtctct cgtaatcacc gagcgaggat 100  
 Leu Cys His Gln  
 20  
 gtagcttgtc atagcttggga cggagacctt ggatgaactc cgtgcgtgcg tgcgtgtgtg 160  
 ggtgtgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtgac ggagaccttg catgaactcc 220  
 gcgtgtgtgt ttgtgtgtgt atctgtatct gtctgtagca gggtctcccg cccctgttaa 280  
 ttatatattt tttctgctgg tttttcattg ctagtctcgt agttcattct aactagaaag 340  
 gtaaaaacttt cttttaataa taccattat aaatgcactt aataatacct ttaattgtgc 400  
 ctttaataat acccaca 417

<210> 25  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

<400> 25  
 cggctcctca tggcgacacg 20

<210> 26  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

<400> 26  
 tgatggcaca ggatggtctg 20

<210> 27  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense Oligonucleotide

<400> 27  
 tagctggctg gaagcaagcc 20

<210> 28  
 <211> 20



<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 28  
ctgtacacat tatctcggac 20

<210> 29  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 29  
gcccaaacc caccagcca 20

<210> 30  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 30  
cttcactaca ctctgctcca 20

<210> 31  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 31  
tgtctgatca tgcagtgcag 20

<210> 32  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 32  
actttatcca cctgtctgat 20

<210> 33  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 33  
tccaactgca ggtgtcccca 20

<210> 34  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 34  
agtcctgagg cagtcatttg 20

<210> 35  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 35  
ctaggctgtg gatgatatgg 20

<210> 36  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 36  
gttttatatg cagcttcaat 20

<210> 37  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 37  
cccacctttc acaccaaaca 20

<210> 38  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 38  
aggacatgga taactgattg 20

<210> 39

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 39  
ccctgaagct tggatgatgat 20

<210> 40  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 40  
aaccataacg accctgaagc 20

<210> 41  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 41  
ttaggagttt tatatccatc 20

<210> 42  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 42  
aatagcttca gctcagctgg 20

<210> 43  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 43  
gtccagaaca atggccattc 20

<210> 44  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 44

tatattcttg aacctgttct

20

<210> 45

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 45

ttgcccttga tgaggactgc

20

<210> 46

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 46

ccccacatgt gaggcaattt

20

<210> 47

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 47

tagagactga cccccacatgt

20

<210> 48

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 48

tccctctgcc atcaagcttc

20

<210> 49

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 49

agaatggaga cttgaaccac

20

<210> 50  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 50  
ttgatttctt ctgtttcagc 20

<210> 51  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 51  
tggctgagaa tacgagctgg 20

<210> 52  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 52  
gtccactgag tttcattcta 20

<210> 53  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 53  
gtgaaagtaa agatagtttt 20

<210> 54  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 54  
aaactgtgga gtgaaagtaa 20

<210> 55  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 55  
tagaactggt gctggtctat 20

<210> 56  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 56  
tccagagcca ggtagaactg 20

<210> 57  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 57  
cagcggctac agaggtagga 20

<210> 58  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 58  
ggtagggctgg cctgtcatcc 20

<210> 59  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 59  
ttccatcttc atcaagcatg 20

<210> 60  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 60  
aggatacttg aattcaagct 20

<210> 61  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 61  
catttttcgg agtgctgcca 20

<210> 62  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 62  
tctgacaatt tacctgtttg 20

<210> 63  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 63  
tgttgtcaaa aactctgaca 20

<210> 64  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 64  
caagatgttg tcaaaaactc 20

<210> 65  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 65  
ggtccagggt ccatgaagct 20

<210> 66  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 66  
gcttaccctc aggtccaggg 20

<210> 67  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 67  
atcataatct tcactgtaca 20

<210> 68  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 68  
ttcatcacca cagcgagcat 20

<210> 69  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 69  
accgatctag ccctcccttc 20

<210> 70  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 70  
accaaggaca ttaagtcgca 20

<210> 71  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 71  
atacatgtga acattctgta 20



<210> 72  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 72  
ttgaggtctc cttcaactgt 20

<210> 73  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 73  
ttccagtcag gtcctttcat 20

<210> 74  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 74  
gtgcaggcct catcaagtgc 20

<210> 75  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 75  
tggaggaagt cctactgtca 20

<210> 76  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 76  
actggcttca tctatcagct 20

<210> 77  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 77  
ccattatttc ctgtgtaagg 20

<210> 78  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 78  
tgccataact tgtatgatca 20

<210> 79  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 79  
tctgtggctt cctcagctga 20

<210> 80  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 80  
gactgagatt catcaggccc 20

<210> 81  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 81  
ctgagaatgt gatgcaggag 20

<210> 82  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 82  
cttcgttcca ctccaaactc 20

<210> 83  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 83  
ttgaatcagt gggacgaacg 20

<210> 84  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 84  
gctccaacag caccaatctc 20

<210> 85  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 85  
tcacttttta actgcatgat 20

<210> 86  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 86  
gaaattccac ctgctttatc 20

<210> 87  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 87  
ttgacgacta tcttttagatg 20

<210> 88  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 88  
gagaatttaa tctcacctgg 20

<210> 89  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 89  
ttcaggacag actccacatg 20

<210> 90  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 90  
acaaggatgg cttcaaccag 20

<210> 91  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 91  
atatctgccg gcattgtgag 20

<210> 92  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 92  
atgaatttca atatctgccg 20

<210> 93  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 93  
tggtcttgaa ggaacaagtc 20

<210> 94  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 94  
aacatggtat catctgcgcc 20

<210> 95  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 95  
atcctttgcc aacatggtat 20

<210> 96  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 96  
agagtacaga tgccagatgc 20

<210> 97  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 97  
gccactgggt gcactgtcat 20

<210> 98  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 98  
aggtaggtca tggcgccaaa 20

<210> 99  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 99  
aactcctgca cgtaggtggc 20

<210> 100  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 100  
ccaaagccct cattgcatgg 20

<210> 101  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 101  
gtccaagcta tgacaagcta 20

<210> 102  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 102  
ataatgggta ttattaaaag 20

<210> 103  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 103  
ccggctgcag aggtaggaga 20

<210> 104  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 104  
tcaccggct gcagaggtag 20

<210> 105  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 105  
tcctccaccg gctgcagagg 20

<210> 106  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 106  
gtcatcctcc accggctgca 20

<210> 107  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 107  
gggctggcct gtcacacctc 20

<210> 108  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 108  
agtgatgggtg ggctggcctg 20

<210> 109  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 109  
aaagtgatgg tgggctggcc 20

<210> 110  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 110  
agcatggtgt gtgagatggg 20

<210> 111  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 111  
cagcatggtg tgtgagatgg 20

<210> 112  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 112  
catccagcat ggtgtgtgag 20

<210> 113  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 113  
cttcatccag catggtgtgt 20

<210> 114  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 114  
tcttcatcca gcatggtgtg 20

<210> 115  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 115  
tccatcttca tccagcatgg 20



<210> 116  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 116  
ggctccatct tcatccagca 20

<210> 117  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 117  
gctggctcca tcttcatcca 20

<210> 118  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 118  
caagctggct ccatcttcat 20

<210> 119  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 119  
ttgaattcaa gctggctcca 20

<210> 120  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 120  
ggatacttga attcaagctg 20

<210> 121  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 121  
caggatactt gaattcaagc 20

<210> 122  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 122  
ccaggatact tgaattcaag 20

<210> 123  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 123  
tgccaggata cttgaattca 20

<210> 124  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 124  
ctgccaggat acttgaattc 20

<210> 125  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 125  
gctgccagga tacttgaatt 20

<210> 126  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 126  
tgctgccagg atacttgaat 20

<210> 127  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 127  
gagtgtgcc aggatacttg 20

<210> 128  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 128  
ggagtgtgc caggatactt 20

<210> 129  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 129  
tccggagtgc tgccaggata 20

<210> 130  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 130  
ttttccggag tgctgccagg 20

<210> 131  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 131  
cattttccgg agtgctgcca 20

<210> 132  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 132  
gcattttccg gaggctgcc 20

<210> 133  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 133  
cctgcatttt ccggagtgc 20

<210> 134  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 134  
catcctgcat tttccggagt 20

<210> 135  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 135  
aaccatcctg cattttccg 20

<210> 136  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 136  
aataaccatc ctgcattttc 20

<210> 137  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 137  
caaaataacc atcctgcatt 20

<210> 138  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 138  
accaaaataa ccatcctgca 20

<210> 139  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 139  
ccaccaaagt aaccatcctg 20

<210> 140  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 140  
ggccccacca aaataacct 20

<210> 141  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 141  
atcctggccc caccaata 20

<210> 142  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 142  
tttggatcct ggccccacca 20

<210> 143  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 143  
cctgtttgga tcctggcccc 20

<210> 144  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 144  
ttacctgttt ggatcctggc 20

<210> 145  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 145  
agcttacctg ttggatcct 20

<210> 146  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 146  
gacagcttac ctgtttggat 20

<210> 147  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 147  
tctgacagct tacctgtttg 20

<210> 148  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 148  
aactctgaca gcttacctgt 20

<210> 149  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 149  
aaaaactctg acagcttacc 20

<210> 150  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 150  
gtcaaaaact ctgacagctt 20

<210> 151  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 151  
gttgtcaaaa actctgacag 20

<210> 152  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 152  
gatgttgtca aaaactctga 20

<210> 153  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 153  
cagcaagatg ttgtcaaaaa 20

<210> 154  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 154  
gtgcagcaag atgttgtcaa .20

<210> 155  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 155  
tgtgtgcagc aagatgttgt 20

<210> 156  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 156  
tgaagcttaa gtgtgtgcag 20

<210> 157  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 157  
gggtccatga agcttaagtg 20

<210> 158  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 158  
tccagggtcc atgaagctta 20

<210> 159  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 159  
taatcttcgc tgtacagctt 20



<210> 160  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 160  
gtcttcatca taatcttcgc 20

<210> 161  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 161  
tagtcttcat cataatcttc 20

<210> 162  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 162  
tcatagtctt catcataatc 20

<210> 163  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 163  
tcatcatcat agtcttcatc 20

<210> 164  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 164  
ccagctcatc atcatagtct 20

<210> 165  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 165  
gtccagctca tcatcatagt 20

<210> 166  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 166  
agtccagctc atcatcatag 20

<210> 167  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 167  
agagtcagc tcatcatcat 20

<210> 168  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 168  
tgccagagtc cagctcatca 20

<210> 169  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 169  
atccagttgc cagagtcag 20

<210> 170  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 170  
catccagttg ccagagtcga 20

<210> 171  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 171  
atccatccag ttgccagagt 20

<210> 172  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 172  
gctatccatc cagttgccag 20

<210> 173  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 173  
catagctatc catccagttg 20

<210> 174  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 174  
gagtcatagc tatccatcca 20

<210> 175  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 175  
cgtgttgagt catagctatc 20

<210> 176  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 176  
acgtgttgag tcatagctat

20

<210> 177  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 177  
cattacgtgt tgagtcata

20

<210> 178  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Antisense Oligonucleotide

<400> 178  
gcattacgtg ttgagtcata

20

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/26608

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/04; A61K 48/00; C12Q 1/68; C12N 15/63  
US CL : 514/44; 435/6, 91.1, 325, 375; 536/23.1, 24.5

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 514/44; 435/6, 91.1, 325, 375; 536/23.1, 24.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
none

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Medline, Embase, Scisearch, Caplus, West

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PHIL-CAREY, K. Isis To Restructure As Crohn's Disease Drug Fails In Phase III. BioWorld Today. 16 December 1999, Vol. 10, No. 239, pages 1 and 2, see entire document.	1-18
A	GEWIRTZ et al. Facilitating oligonucleotide delivery: Helping antisense deliver on its promise. Proc. Natl. Acad. Sci. April 1996, vol. 93, pages 3161-3163, see entire document.	1-18
A	BRANCH, A. A good antisense molecule is hard to find. TIBS. February 1998, vol. 23, pages 45-50, see entire document.	1-18

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance  
"B" earlier application or patent published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
"&" document member of the same patent family

Date of the actual completion of the international search

08 October 2001 (08.10.2001)

Date of mailing of the international search report

30 OCT 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Andrew Wang

Telephone No. 703-308-0196